


Journal Pre-proof

Photodegradation of Organic Micropollutants in Aquatic Environment: Importance, Factors and Processes

Zhongyu Guo , Dilini Kodikara , Luthfia Shofi Albi , Yuta Hatano , Guo Chen , Chihiro Yoshimura , Jieqiong Wang

PII: S0043-1354(22)00199-3
DOI: <https://doi.org/10.1016/j.watres.2022.118236>
Reference: WR 118236



To appear in: *Water Research*

Received date: 29 October 2021
Revised date: 22 February 2022
Accepted date: 23 February 2022

Please cite this article as: Zhongyu Guo , Dilini Kodikara , Luthfia Shofi Albi , Yuta Hatano , Guo Chen , Chihiro Yoshimura , Jieqiong Wang , Photodegradation of Organic Micropollutants in Aquatic Environment: Importance, Factors and Processes, *Water Research* (2022), doi: <https://doi.org/10.1016/j.watres.2022.118236>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2022 Published by Elsevier Ltd.

**Photodegradation of Organic Micropollutants in Aquatic Environment:
Importance, Factors and Processes**

Zhongyu Guo^a, Dilini Kodikara^a, Luthfia Shofi Albi^a, Yuta Hatano^a, Guo Chen^a,
Chihiro Yoshimura^{a,*}, Jieqiong Wang^b

a Department of Civil and Environmental Engineering, Tokyo Institute of Technology,
Meguro-ku, Tokyo, 152-8552, Japan

b Research Center for Eco-environmental Engineering, Dongguan University of
Technology, Dongguan 523808, China


*Corresponding author

Chihiro Yoshimura, Phone: 03-5734-3548; Email: yoshimura.c.aa@m.titech.ac.jp

Highlights

- Importance of photodegradation in removal of OMPs in waters was discussed.
- Processes of OMPs' photodegradation was compared to biodegradation.
- Understanding on roles of DOM on photodegradation is concisely updated.
- Major knowledge gaps in environmental photochemistry are identified.

Graphical Abstract



Abstract

Photochemical reactions widely occur in the aquatic environment and play fundamental roles in aquatic ecosystems. In particular, solar-induced photodegradation is efficient for many organic micropollutants (OMPs), especially those that cannot undergo hydrolysis or biodegradation, and thus can mitigate chemical pollution. Recent reports indicate that photodegradation may play a more important role than biodegradation in many OMP transformations in the aquatic environment. Photodegradation can be influenced by the water matrix such as pH, inorganic ions, and dissolved organic matter (DOM). The effect of the water matrix such as DOM on photodegradation is complex, and new insights concerning the disparate effects of DOM have recently been reported. In addition, the photodegradation process is also influenced by physical factors such as latitude, water

depth, and temporal variations in sunlight as these factors determine the light conditions. However, it remains challenging to gain an overview of the importance of photodegradation in the aquatic environment because the reactions involved are diverse and complex. Therefore, this review provides a concise summary of the importance of photodegradation and the major processes related to photodegradation of OMPs, with particular attention given to recent progress on the major reactions of DOM. In addition, major knowledge gaps in this field of environmental photochemistry are highlighted.

Keywords: Reactive intermediates, water matrix, dissolved organic matter, photodegradation, physical factors.

1. Introduction

Photochemistry is deeply integrated into the Earth's biological and ecological systems. In the atmosphere, the presence of oxygen and its photochemical product, ozone, screen out most ultraviolet radiation (UV) from the sun, making life on earth possible. Visible light, which is transmitted through the atmosphere, is intimately involved in various biological and chemical phenomena, such as vision, photosynthesis, and photodegradation. Those phenomena are explained by the sequential reactions caused by excited states of atoms and molecules (Wayne, 2005). As such, solar light and the induced photochemical processes are fundamental aspects of ecosystems and the environment. Consequently, photochemistry often appears as a dominant process in health and environmental problems as well. Nucleic acids and proteins are readily damaged by short-wavelength radiation (i.e., UV) and ecosystems can be disrupted by excess or unwanted light (i.e., light pollution). Except for depletion of the ozone layer, such alterations of the light environment have been investigated to date mostly in relation to plant and animal physiology, but not aquatic photochemistry and microbiology.

Organic micropollutants (OMPs) are a serious concern due to their occurrence, persistence, and toxicity in aquatic environments. For instance, pharmaceuticals and personal care products (PPCPs), agrochemicals, dyes, fire retardants, UV filters, and antioxidants have been frequently detected and worldwide in many water bodies including coasts, rivers, and lakes (Madikizela et al., 2020). Given their unexpected interaction with organisms, a certain portion of OMPs is detrimental to aquatic organisms (Rozas et al., 2017), including bacteria, algae, invertebrates (Brausch and Rand, 2011; Hernando et al., 2006; Michael et al., 2014) and fish (Crane et al., 2006; Schwaiger et al., 2004). The direct consequence of antibiotic residues is the

development of antibiotic-resistant bacterial strains (Grenni et al., 2018), which can potentially cause severe disease, including urinary tract, lower respiratory tract, and bloodstream infections (Sanganyado and Gwenzi, 2019). In addition, even at low concentrations, long-term exposure to OMPs increases the ecological risk to aquatic organisms in the water environment (Kumar et al., 2019; Yang et al., 2020). Generally in aquatic ecosystems, each OMP can exhibit a wide range of concentrations (ng/L to $\mu\text{g/L}$ level) (Kameda et al., 2011; Patel et al., 2019; Petrović et al., 2014; Vimalkumar et al., 2018; Zareitalabad et al., 2013). Relatively high OMP concentrations have commonly been observed in water bodies that receive treated wastewater because conventional wastewater treatment plants are not equipped to eliminate OMPs efficiently (Gruchlik et al., 2018; Knopp et al., 2016; Margot et al., 2013; Ribeiro et al., 2015; Söregård et al., 2019).

Photodegradation plays an essential role in the fate of OMPs, particularly in the surface layer of water bodies, where UV and visible light are abundant during the daytime, inducing direct and indirect photolysis of OMPs. In direct photolysis, OMPs undergo various photodegradation paths depending on their chemical structure, and the direct absorption of solar radiation is enhanced by the presence of light-absorbing functional groups such as aromatic rings (Boreen et al., 2003; Challis et al., 2013; Edlund et al., 2006; Fatta-Kassinos et al., 2011). Indirect photodegradation in natural water involves photochemically produced reactive intermediates (PPRIs) such as singlet oxygen ($^1\text{O}_2$) and hydroxyl radical ($\text{HO}\cdot$), which can initiate photodegradation of OMPs (Vione et al., 2014). Overall, the rate of photodegradation depends on the chemical structure of the OMPs, the irradiance (determined by the photon flux in a particular wavelength band), and the water constituents present (e.g., DOM; Boreen et al., 2004; Fent et al., 2006).

A growing number of articles on such underwater photolysis have been published. However, a majority of these articles describe processes and mechanisms under well-controlled conditions and thus it is often difficult to place such knowledge in the context of photodegradation of OMPs in the actual aquatic environment. Consequently, the importance of photodegradation in the real environment is largely unknown compared with other pathways such as biodegradation. In fact, several articles describe *in situ* photochemical process (Bouhala et al., 2020; Dimou et al., 2005; Shih and Wang, 2009). Nevertheless, these articles are still scarce and fragmented in terms of irradiation conditions, processes, and target OMPs, and thus comparative studies (e.g., Baena-Nogueras et al., 2017; Ossola et al., 2021) are required to obtain a comprehensive overview of photodegradation of OMPs in the aquatic environment in order for us to undertake a quantitative environmental assessment of OMPs.

It is widely accepted that the photodegradation of OMPs largely depends on the water matrix as the water matrix determines the production of PPRIs as well as inhibitors of photolysis (Challis et al., 2014; McNeill and Canonica, 2016; Sharpless and Blough, 2014; Vione et al., 2014; Vione and Scozzaro, 2019; Yan and Song, 2014). Nevertheless, knowledge of photodegradation processes in a wide variety of water matrices including the complex photoreactions of DOM is still too limited to obtain a comprehensive overview. In addition, photodegradation is also affected by physical environmental factors such as water depth, latitude, season, and time of day. These physical factors have received less attention than the water matrix although their impacts need to be elucidated as a basis for investigating the fate of OMPs in the aquatic environment and as for enhancing knowledge of environmental photochemistry in the broader sense.

Given these environmental processes that have yet to be revealed, we believe that it would be useful to update the existing knowledge of the field in order to achieve more effective management of aquatic photodegradation of OMPs. This review, therefore, aims to provide a concise overview of the importance of photochemical reactions in the fate of OMPs in the aquatic environment from a comprehensive viewpoint and addresses the basics of aquatic photochemistry and its relation to environmental factors including the water matrix and physical conditions.

2. Why photodegradation?

Solar energy is abundant on the surface layer of water bodies. Despite atmospheric absorption and scattering, about 72% of the solar energy reaches the Earth's surface (Valero et al., 2000), but the short wavelengths (≤ 280 nm) are completely absorbed. Consequently, direct solar irradiance is decreased from 1361 W m^{-2} at the top of the atmosphere to about $1,000 \text{ W m}^{-2}$ at the Earth's surface (Coddington et al., 2016; Myers, 2013). The total energy that reaches the Earth's surface is about 3×10^{24} J per year, which is equivalent to about 6500 times the annual energy consumption of humankind (4.6×10^{20} J per year) (Klan and Wirz, 2009). Within a water body, OMPs are mainly degraded and transformed by biological (e.g., biodegradation) and chemical (e.g., photolysis, hydrolysis) processes. Among them, hydrolysis is usually slow and requires the existence of hydrolytic functional groups in the OMP such as ester bonds (Xu et al., 2019, 2021) while photodegradation and biodegradation occur more extensively in the water environment. Owing to the abundant solar radiation and various photosensitizers (i.e., DOM) in the water, photodegradation may be a pivotal removal pathway for many OMPs, with possible interactions with other processes.

Baena-Nogueras et al. (2017) have reported that the photodegradation and biodegradation rate constants of 33 frequently detected PPCPs were significantly larger than their hydrolysis rates. Furthermore, most of the studied PPCPs exhibited faster photodegradation (half-life, $t_{1/2}$ range 0.02–29 h) than biodegradation ($t_{1/2}$ range 4–2310 h) (Figure 1), while the water matrix (pure water, freshwater, and seawater) affects the rate. In addition, typical PPCPs such as ibuprofen (IBP), diclofenac (DCF) and sulfamethoxazole (SMX), respectively, exhibited photodegradation rates of 0.03 h^{-1} ($t_{1/2}$ 23 h), 4.04 h^{-1} ($t_{1/2}$ 0.17 h), and 0.19 h^{-1} ($t_{1/2}$ 3.73 h) in spiked pure water under simulated solar light. In contrast, IBP exhibited a biodegradation rate of 0.0005 h^{-1} ($t_{1/2}$ 1386 h) in spiked freshwater, while DCF and SMX exhibited no biodegradation at all for 28 days (Baena-Nogueras et al., 2017). These studies clearly indicated that photodegradation may play a more important role in many OMPs transformation than biodegradation.

On the other hand, many OMPs are toxic to microorganisms (Ebert et al., 2011; Larsson et al., 2007; Le Page et al., 2017; Patel et al., 2019; Rubasinghege et al., 2018) partly because of the intended uses of the OMPs, for example, as antimicrobial agents. This toxicity poses significant challenges in terms of biodegradability in the environment. Hence, photodegradation out of all the potential degradation and transformation processes possibly plays a rather important role in deciding the fate of OMPs in aquatic environments. It is worth noting that photochemical transformation of OMPs, including photodegradation in some cases, may result in photo-induced toxicity and generate more persistent products, which possibly increases ecological risk (Buth et al., 2007; Latch et al., 2003a; Plumlee et al., 2009; Robert et al., 2017). For example, the photodegradation of fenoxaprop-*p*-ethyl forms products with higher toxicity and more persistence (Lin et al., 2008). Simulated solar irradiation can

enhance the acute toxicity of fluoroquinolones and anthraquinones to *Daphnia magna* (Luo et al., 2018; Wang et al., 2009).

3. Overview of aquatic photodegradation

Photodegradation processes can be either direct or indirect (Wang et al., 2017; Zhang et al., 2017; De la Cruz et al., 2013) and Figure 2 shows the overview of aquatic photodegradation. Direct photolysis occurs when OMPs directly receive photon energy from an irradiation source and the OMP's electrons are excited to a higher energy level. This allows the molecule to undergo a chemical transformation, ultimately causing the OMP to photodegrade. An OMP's capability to undergo direct photolysis depends on two factors: the ability of the OMP to absorb photon energy as indicated by its molar absorption coefficient and the likelihood of the molecule to undergo chemical transformation as indicated by its quantum yield (Ge et al., 2009; Yao et al., 2013). Direct photolysis can be influenced by the water matrix via processes such as light screening caused by light-absorbing constituents as well as back reduction by DOM (Liu et al., 2020).

In indirect photodegradation, photons from an irradiation source are received by photosensitizers to produce PPRIs, which ultimately degrade the target OMPs. Therefore, the composition of the water matrix in which the target OMPs occur can have a positive or negative influence on the efficiency of photodegradation (Lado Ribeiro et al., 2019). Indirect photodegradation is largely dependent on water matrix components such as $\text{NO}_3^-/\text{NO}_2^-$ and DOM because these components can photo-generate PPRIs to degrade the OMPs (Ge et al., 2009). However, the impact of such photosensitizers can vary depending on the target OMPs (Wang et al., 2017; Zhang et al., 2017; Sun et al., 2014) and also on the source of DOM (Sun et al., 2014) which will be elaborated on in the next section of this review.

The photodegradation process is active only during daytime when light is abundant and the corresponding rate of photodegradation is highly affected by the light intensity imposed on the OMPs or photosensitizers. In addition, the extent of light attenuation increases with the depth of the water column and varies seasonally (Mathon et al., 2019). Against this backdrop, the influence of natural solar light on the degradation of OMPs remains largely unknown and has been scarcely studied despite that such information is essential for understanding the attenuation process of OMPs and elucidating their fate in the environment (Wang et al., 2021b).

Photodegradation studies are usually carried out under well-controlled conditions in the laboratory and extrapolation of the rate of photodegradation from the laboratory to different environments is of great practical significance for management of chemicals (precursors of OMPs). However, this strategy poses a number of challenges because laboratory conditions are different from those in real water bodies (Mackay, 2001). In recent years, a model called APEX (Aqueous Photochemistry of Environmentally-occurring Xenobiotics) has been developed to simulate the photochemical reactions of aqueous OMPs (Bodrato and Vione, 2014; Vione, 2020). This model is kinetics-based and user friendly. Zhou et al. (2018) have also developed a kinetics-based photodegradation rate prediction model, which takes variations in the intensity of underwater and hourly sunlight into consideration. These two models are useful in extrapolating photochemical persistence data determined in the laboratory to the real environment, although further development is needed so that these models can be used to comprehensively study photodegradation in actual water environments.

With respect to above models, the input photochemical parameters of OMPs such as quantum yield of direct photolysis and the second-order reaction rate constants are necessary for predicting photochemical persistence. However, these

parameters are not available for many OMPs, let alone the thousands of chemicals that have been synthesized and used, and have entered the aquatic environment. Prediction of those photochemical parameters for OMPs is pertinent to persistence evaluation. Some pioneering studies employed quantitative structure-activity/property relationship (QSAR/QSPR) to predict these parameters including the quantum yield for OMPs (Chen et al., 2000; Chen et al., 2001a; Chen et al., 2001b; Niu et al., 2005; Shao et al., 2010; Yang et al., 2003) and the photochemical kinetics parameters (Li et al., 2014; Luo et al., 2017; Luo et al., 2019), which provide an opportunity for prediction given the lack of relevant data.

Nevertheless, this article cannot cover all the basics of environmental photochemistry and so we refer interested readers to relevant texts for further information about the basic mechanisms (Klan and Wirz, 2009; Boule, 1999; Calza and Vione 2015).

4. Effects of the water matrix on photodegradation

Photodegradation of OMPs can be influenced by the constituents in the water matrix such as ions ($\text{NO}_3^-/\text{NO}_2^-$, $\text{Fe}^{2+}/\text{Fe}^{3+}$, $\text{CO}_3^{2-}/\text{HCO}_3^-$, halide ions, etc.), DOM, and suspended solids (Ge et al., 2010a; Ge et al., 2010b; Ge et al., 2010c; Vione et al., 2014). Likewise, coexisting OMPs can be considered as water matrix components that can influence photodegradation. The water matrix can either accelerate photodegradation or inhibit it. In addition, the pH of the water can influence the charge state of DOM and the presence of ionized OMPs in water, meaning that photodegradation is dependent on the pH of the water (Boreen et al., 2004; Ge et al., 2010a; Wei et al., 2013; Werner et al., 2006). For example, ciprofloxacin (CIP) had five dissociated species and each of them had dissimilar photolytic kinetics and

products (Wei et al., 2013). In the following sections, the effects of the water matrix on photodegradation of OMPs are further elaborated.

4.1 Inorganic ions

Inorganic ions are ubiquitous in the aquatic environment and play various roles in the photodegradation of OMPs. The ions can interact with OMPs, altering the photodegradation behaviors of the OMPs. For example, coordination with transition metals such as Cu(II) can alter the light absorption characteristic and photo-reactivity of OMPs (Wei et al., 2015). Some inorganic ions can additionally act as direct/indirect sources of PPRIs, which can provide additional opportunity for degradation of refractory OMPs. Hence, in this section, focus is given mainly to the effects of inorganic ions that can act as sources of PPRIs for the photodegradation of OMPs.


$\text{NO}_3^-/\text{NO}_2^-$ play important roles in the photodegradation of OMPs because $\text{NO}_3^-/\text{NO}_2^-$ are important sources of $\text{HO}\cdot$ and these species are ubiquitous in natural waters. $\text{NO}_3^-/\text{NO}_2^-$ can produce $\text{HO}\cdot$ through direct photolysis under the action of solar radiation ($\lambda > 280 \text{ nm}$); the main reactions are shown below (Mack and Bolton, 1999; Wu et al., 2020):





As shown in Eq. (1)-(3), both NO_3^- and NO_2^- can directly absorb solar radiation and produce $\text{O}^- \cdot$, which is a precursor of $\text{HO}\cdot$. The pK_a of $\text{HO}\cdot$ is 11.9 (eq.3), which means that at environmental pH, reaction (3) favors the generation of $\text{HO}\cdot$ (Mack and Bolton, 1999). NO_2^- can be transformed to its conjugated acid HONO which is more efficient in producing $\text{HO}\cdot$, thus increasing the production of $\text{HO}\cdot$ with decreasing the

pH. However, this pH effect is clearly seen only in water with pH < 5, which is not a common pH in surface waters (Arakaki et al., 1999). NO₃⁻ can photo-isomerize into OONO⁻ and form its conjugated acid HOONO with pKa ≈ 6.5-7, where HOONO can produce HO· while OONO⁻ cannot (Gligorovski et al., 2015; Mack and Bolton, 1999). Nevertheless, this pathway may be also insignificant in the natural environment because the quantum yield for photoproduction of OONO⁻ under solar radiation is low (less than 0.26% at 313 nm irradiation) (Benedict et al., 2017). Although the concentration of NO₂⁻ is usually lower than that of NO₃⁻ in natural waters, NO₂⁻ absorbs solar radiation more effectively and produces HO· with a higher quantum yield ($\Phi_{\text{NO}_2}^{\text{HO}\cdot} = 0.025\text{-}0.065$, depending on wavelength; $\Phi_{\text{NO}_3}^{\text{HO}\cdot} \approx 0.01$) (Vione et al., 2014). Consequently, under the typical ratio for [NO₃⁻]/[NO₂⁻] which occurs in natural waters, NO₂⁻ is more efficient than NO₃⁻ in producing HO· (two orders of magnitude more efficient) (Gligorovski et al., 2015; Minero et al., 2007).

Fe ion species, particularly Fe(II)/Fe(III) complexes (mainly hydroxo and organic complexes), can participate in Fenton/Photo-Fenton reactions to generate HO· that causes OMPs degradation in sunlit aquatic environments. The photochemistry of Fe has been reviewed previously, and two representative equations (Eq. (4) and (5)) concerning HO· production are shown below (Lueder et al., 2020; Vione et al., 2014). The importance of these reactions depends on specific water bodies where the pH is low enough and the concentrations of Fe(II)/Fe(III) are sufficiently high (Gligorovski et al., 2015). This is because Fenton/Photo-Fenton reactions are favored under low pH condition (pH = 2~3) (Feng and Nansheng, 2000).





Photodegradation of OMPs can be influenced by $\text{CO}_3^{2-}/\text{HCO}_3^-$, mainly due to the generation of the carbonate radical ion ($\text{CO}_3^{\cdot-}$) that reacts with OMPs in sunlit water bodies. $\text{CO}_3^{\cdot-}$ is a selective oxidant with $E_{\text{NHE}}(\text{CO}_3^{\cdot-}/\text{CO}_3^{2-}) = +1.59 \text{ V}$ (Huie et al., 1991), which is lower than that of $\text{HO}\cdot$ while its steady-state concentration is estimated to be two orders of magnitude higher than that of $\text{HO}\cdot$ (Sulzberger et al., 1997). The reactions between $\text{CO}_3^{\cdot-}$ and OMPs mainly occur via electron transfer or a hydrogen abstraction mechanism (Liu et al., 2018). Thus, $\text{CO}_3^{\cdot-}$ plays an important role in photodegradation of some OMPs containing electron-rich structures such as nitrogen- and sulfur-containing compounds (thiols), and aromatic compounds (anilines and phenols) (Busset et al., 2007; De Laurentiis et al., 2014; Mazellier et al., 2007; Sun et al., 2020; Wu and Linden, 2010). $\text{CO}_3^{\cdot-}$ is a secondary radical because, as shown in Eq. (6) and (7), the $\text{CO}_3^{\cdot-}$ is generated through either reaction between $\text{CO}_3^{2-}/\text{HCO}_3^-$ and $\text{HO}\cdot$ or between CO_3^{2-} and excited species (i.e., excited triplet state of DOM and OMPs ($^3\text{DOM}^*/^3\text{OMPs}^*$)) (Yan et al., 2019).



Reaction (6) is highly pH-dependent because of the dissociation of carbonic acid to CO_3^{2-} and HCO_3^- occurs at different pH values ($\text{p}K_{\text{a}1} = 6.3$ and $\text{p}K_{\text{a}2} = 10.3$) (Vione et al., 2014). The generation of $\text{CO}_3^{\cdot-}$ is more strongly favored when the pH increases from 6 to 9 (circumneutral pH) as CO_3^{2-} is more reactive than HCO_3^- toward $\text{HO}\cdot$ ($k_{\text{HCO}_3^-, \text{HO}\cdot} = 8.5 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$, $k_{\text{CO}_3^{2-}, \text{HO}\cdot} = 3.9 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$), and the proportion of CO_3^{2-} increases when the pH increases (Zhou et al., 2020). Eq. (7) is important only under specific conditions (waters with high DOC) (Canonica et al., 2005) and in most cases, Eq. (6) is the dominant reaction for generating $\text{CO}_3^{\cdot-}$ in natural water bodies.

Halide ions can react mainly react through an oxidation reaction with photochemically produced HO· or ³DOM* to form various halogen radical species (X·, X₂⁻· and XY· where X and Y stand for Cl, Br, or I) (Parker and Mitch, 2016; Zhang and Parker, 2018; Zhang et al., 2018a):



Compared with X·, X₂⁻· and XY· are dominant in halide-containing sunlit waters (concentrations are 3 orders of magnitude higher than X·) because halide ions are abundant (Yang and Pignatello, 2017). In seawater, the steady-state concentrations of Br· and BrCl· are estimated to be 3–4 orders of magnitude higher than the concentration of HO· (Zhang and Parker, 2018). These halogen radicals can be quenched by HCO₃⁻/CO₃⁻ to form CO₃⁻· and by DOM to form DOM radical. Halogen radicals can react with OMPs via halogenation (Cheng et al. 2021), and many polyhalogenated OMPs can be produced; these OMPs can exhibit serious ecological risks (Lallas, 2001). On the other hand, halogen radicals can also react with OMPs through the oxidation reactions where halogen atoms are not introduced into the reaction products (Wang et al. 2021). Since the reaction between halogen radicals and the OMPs is a redox reaction, the redox potential is pertinent to the reaction. The reduction potential of Cl· ($E_{\text{NHE}} = 2.5 \text{ V}$) is the highest among all the halogen radicals (others range from 1.1 to 2.2 V) and is slightly lower than that of HO· ($E_{\text{NHE}} = 2.7 \text{ V}$),

indicating that $\text{Cl}\cdot$ has high reactivity toward many OMPs (Wardman, 1989). Halogen radicals can react with saturated aliphatic compounds (via hydrogen abstraction), olefinic compounds (via addition reaction), aromatic compounds (via addition reaction, electron abstraction), and organosulfur species (adduct formation or sulfur oxidation) (Wang et al., 2021a; Zhang and Parker, 2018).

In addition to the generation of halogen radicals, Parker et al. (2013) have reported that an increase in ionic strength can slow down intra-organic matter electron transfer of $^3\text{DOM}^*$, resulting in higher steady-state concentrations of $^3\text{DOM}^*$ ($[^3\text{DOM}^*]_{\text{ss}}$). Elevation of $[^3\text{DOM}^*]_{\text{ss}}$ by halide ions has also been observed in further studies as well as the quenching of excited singlet-state DOM ($^1\text{DOM}^*$) (Glover and Rosario-Ortiz, 2013). Higher $[^3\text{DOM}^*]_{\text{ss}}$ can promote photodegradation when $^3\text{DOM}^*$ is critical to the photodegradation (Chen et al. 2019; Li et al. 2016). Suppression of photodegradation by halide ions was also observed for 17β -estradiol, a phenol, and this was attributed to the inhibition of electron transfer pathways between $^3\text{DOM}^*$ and 17β -estradiol (Grebel et al. 2012; Parker et al. 2013). Further research is needed to clarify the relative importance between elevation of $[^3\text{DOM}^*]_{\text{ss}}$ and inhibition of the electron transfer pathways by halide ions in the photodegradation of OMPs. Regarding halide-specific effects, previous studies have found that Cl^- and Br^- can enhance photobleaching of DOM (Grebel et al. 2009; Grebel et al. 2012), thus decreasing the sensitizing ability of DOM and inhibiting photodegradation. Halide ions like Br^- may also exhibit a “heavy atom” effect on photodegradation, which can promote intersystem crossing from the excited singlet state to the excited triplet state and influence the photodegradation (Koziar and Cowan, 1977; Li et al. 2016).

4.2 Dissolved organic matter

DOM is ubiquitous in environmental water bodies, and it can widely absorb solar radiation up to 500 nm (Nebbioso and Piccolo, 2013). After absorbing light, DOM can produce many PPRIs such as $^1\text{DOM}^*$, aqueous electron (e^-_{aq}), superoxide ($\text{O}_2^{\cdot-}$), H_2O_2 , $^3\text{DOM}^*$, $^1\text{O}_2$, and $\text{HO}\cdot$, which contribute to photodegradation (Figure 3). DOM can also inhibit photodegradation, for example, by competing for light with OMPs and quenching PPRIs generated during photodegradation.

4.2.1 Promotion effects

Carbonyl structures and aromatic compounds are the main chromophores of DOM. These chromophores absorb solar radiation and first produce $^1\text{DOM}^*$ with a short lifetime and low steady-state concentration in environmental waters (Boyle et al., 2009). However, $^1\text{DOM}^*$ has high reactivity and has been shown to play a pivotal role in the formation of some PPRIs such as e^-_{aq} and charge-transfer-state DOM ($\text{DOM}^{+/-\cdot}$).

The excited singlet-state aromatic groups, mainly phenol and its derivatives (such as aromatic carboxylic acids (Köhler et al., 1985) and aromatic amino acids (Mossoba et al., 1982)) in $^1\text{DOM}^*$, can eject e^-_{aq} . The e^-_{aq} generated plays an insignificant role in OMPs attenuation in environmental waters because it is produced only inside the DOM microenvironment and mostly recombines with cations before escaping into the bulk solution (Zepp et al., 1987). Only very hydrophobic OMPs with a high octanol-water partition coefficient K_{OW} , such as Mirex ($K_{\text{OW}} \approx 10^{6.29}$; Burns et al., 1996, 1997), can bind with DOM so that e^-_{aq} can react to them.

$^1\text{DOM}^*$ can also produce $\text{DOM}^{+/-\cdot}$ through direct intramolecular electron transfer (McKay et al., 2016; Sharpless and Blough, 2014). These $\text{DOM}^{+/-\cdot}$ states are formed as a result of the transfer of an electron either partially or completely from an electron-rich donor (e.g., hydroxy- or alkoxy-benzenes) to an electron-poor acceptor

(e.g., carbonyl-containing moieties like aromatic ketones and quinones) (McKay et al., 2016). Up to now, $\text{DOM}^{+/-}$ is regarded as the major PPRI responsible for the production of O_2^- (Ma et al., 2020; Zhang et al., 2012). Electron-rich phenolic moieties are precursors of $\text{DOM}^{+/-}$, which further verified that $\text{DOM}^{+/-}$ is the precursor of O_2^- . O_2^- often acts as a weak oxidant with $E_{\text{NHE}}(\text{O}_2^-/\text{H}_2\text{O}_2) = +0.91 \text{ V}$ and it plays an important role in redox speciation of dissolved metals in natural water bodies (Ma et al., 2020). Catalyzed or uncatalyzed disproportionation of O_2^- can produce H_2O_2 (Cooper and Lean, 1989; Petasne and Zika, 1987; Scully et al., 1996), which is an important PPRI because it can further generate $\text{HO}\cdot$ through Fenton reactions (Southworth and Voelker, 2003). Although only about 12-35% of O_2^- is transformed to H_2O_2 via dismutation (Le Roux et al., 2021) and another portion of O_2^- underwent photoinduced loss possibly by reacting with DOM^+ (Ma et al., 2020), O_2^- is still considered the most important PPRI for producing H_2O_2 .

Aromatic carbonyl and quinones groups in $^1\text{DOM}^*$ (about 4%-8%) can be transformed to $^3\text{DOM}^*$ through intersystem crossing (McKay, 2020). Although $^3\text{DOM}^*$ is less reactive than $^1\text{DOM}^*$, $^3\text{DOM}^*$ has a longer lifetime and higher steady-state concentration (10^{-14} to 10^{-12} M), making $^3\text{DOM}^*$ an important PPRI in environmental waters (McNeill and Canonica, 2016). $^3\text{DOM}^*$ can directly react with OMPs through oxidation (electron transfer, proton-coupled electron transfer, or other related mechanisms) or energy transfer (McNeill and Canonica, 2016). $^3\text{DOM}^*$ is also the precursor of $^1\text{O}_2$ and $\text{HO}\cdot$ (mainly $\text{HO}\cdot$ -like species).

$^3\text{DOM}^*$ tends to react with aromatic anilines and electron-rich phenols via an oxidation mechanism and $^3\text{DOM}^*$ mainly acts as an oxidant, which should have a higher redox potential than the target compound. Plenty of examples of oxidation with $^3\text{DOM}^*$ were summarized by McNeill and Canonica (2016). For energy transfer

reactions with $^3\text{DOM}^*$, a prerequisite is that the compound should have a lower excited triplet energy (E_T) than $^3\text{DOM}^*$. The reaction between O_2 and $^3\text{DOM}^*$ forming $^1\text{O}_2$ is a typical energy transfer reaction (Zepp et al., 1977). $^1\text{O}_2$ occurs in sunlit DOM solutions with steady-state concentrations ranging from 10^{-12} to 10^{-15} M, which is in the same range of $^3\text{DOM}^*$ (McNeill and Canonica, 2016; Peterson et al., 2012). $^1\text{O}_2$ is a selective oxidant and can react rapidly with only certain OMPs such as phenolates and furans (Ossola et al., 2019) and indoles, imidazoles, and some easily oxidized amino acids (Boreen et al., 2008; Latch et al., 2003b). It is notable that the distributions of $^1\text{O}_2$ exhibit a microheterogeneity because the species is produced in a microenvironment of DOM, and sometimes it reaches higher steady-state concentration in the hydrophobic sites of DOM compared with that in the bulk solution (Grandbois et al., 2008; Latch and McNeill, 2006). Therefore, hydrophobic OMPs that can bind to DOM are possibly susceptible for reaction with $^1\text{O}_2$.

In addition to $^1\text{O}_2$, it is often reported that only compounds containing dienes can react with $^3\text{DOM}^*$ through energy transfer via compounds such as sorbic acid (Grebel et al., 2011), isoprene, and sorbic alcohol (Zhou et al., 2019a) and the corresponding products are their cis-/trans-isomers. $^3\text{DOM}^*$ are mixtures of different excited triplet-state species, which means the E_T is a distribution rather than a definite value.

Zepp et al. (1985) employed diene probes and demonstrated that about 37% of $^3\text{DOM}^*$ had high E_T ($\geq 250 \text{ kJ mol}^{-1}$), which was labeled as high-energy $^3\text{DOM}^*$ (Hi- $^3\text{DOM}^*$). The remainder was labeled as low-energy $^3\text{DOM}^*$ ($94 \text{ kJ mol}^{-1} \leq E_T \leq 250 \text{ kJ mol}^{-1}$). Recently, Zhou et al. (2019a) used sorbic alcohol ($E_T \approx 250 \text{ kJ mol}^{-1}$) as a Hi- $^3\text{DOM}^*$ probe and found that the average proportion of Hi- $^3\text{DOM}^*$ in 15 different types of DOM was 33%, which was very close to the 37% reported previously (Zepp et al., 1985). Notably, the proportion of Hi- $^3\text{DOM}^*$ depended on the source of DOM,

with the proportion being higher in autochthonous freshwater DOM than allochthonous freshwater DOM (Zhou et al., 2019a).

Meanwhile, photodegradation was found to be faster in maricultural seawater DOM than seawater DOM with less mariculture influence, which was attributed to a higher content of $\text{Hi-}^3\text{DOM}^*$ (Guo et al., 2021). Wang et al. (2020a) have also confirmed the high reactivity of $\text{Hi-}^3\text{DOM}^*$ in the photodegradation of OMPs using 11 types of DOMs. However, it is still not clear whether $\text{Hi-}^3\text{DOM}^*$ plays an important role in the overall photodegradation of OMPs in the natural environment as has been assumed.

$^3\text{DOM}^*$ is a precursor of $\text{HO}\cdot$, which is a highly reactive radical that reacts with OMPs at a nearly diffusion controlled rate. $\text{HO}\cdot$ production by DOM also exhibits microheterogeneity similar to that of $^1\text{O}_2$, where more $\text{HO}\cdot$ is possibly produced inside the hydrophobic core of DOM (Yan et al., 2021). Hence, hydrophobic OMPs that can enter the DOM microenvironment are possibly specifically susceptible to photodegradation. However, in sunlit surface waters, the steady-state concentration of $\text{HO}\cdot$ (10^{-16} – 10^{-18} M) is much lower than that of $^3\text{DOM}^*$ or $^1\text{O}_2$ even though direct photolysis of nitrite/nitrate or photo-Fenton reactions can elevate the steady-state concentration of $\text{HO}\cdot$ (Mack and Bolton, 1999; White et al., 2003; Zepp et al., 1992). Regarding the production of $\text{HO}\cdot$, there is still a dispute as to whether $\text{HO}\cdot$ is produced by $^3\text{DOM}^*$ or other precursors like $\text{DOM}^{+/-}\cdot$ in DOM solutions exposed to sunlight (McKay and Rosario-Ortiz, 2015). In addition to free $\text{HO}\cdot$, the generation of $\text{HO}\cdot$ -like species has also been reported, including lower-energy hydroxylation intermediates that react with OMPs at a slightly lower rate than free $\text{HO}\cdot$ reacts with them, and quinones are regarded as the precursors of these species (Gan et al., 2008; Liu et al., 2019; Page et al., 2011). Although studies usually do not distinguish

between them in measurement, the presence of HO·-like species can be confirmed by using methane, which selectively quenches HO· rather than HO·-like species (Page et al., 2011).

Two pathways were proposed for production of HO· namely H₂O₂-dependent and H₂O₂-independent pathways (McKay and Rosario-Ortiz, 2015) while the percentage for the H₂O₂-dependent pathway varied among the DOM samples (Page et al., 2011). Regarding the H₂O₂-independent pathway, Glover and Rosario-Ortiz (2013) found that after quenching of ¹DOM*, the steady-state concentrations of ³DOM* and ¹O₂ both increased while the quantum yield of HO· decreased. This means that HO· and ¹O₂ do not share the same precursor (³DOM*) and that ¹DOM* may contribute to HO· production. Sun et al. (2015) found that direct photolysis of substituted phenols, specifically carboxylate-substituted phenols, may be an important source of HO· in DOM. Two feasible pathways were proposed: (1) the carboxylate-substituted phenol generates an e_{aq}⁻ and a phenol cation radical. Then, the phenol cation radical either undergoes deprotonation and forms HO· or (2) is transformed to a dihydroxycyclohexadienyl radical intermediate that finally eliminates HO·. However, the importance of this mechanism is not clear because good correlation has not been observed between the quantum yield for HO· production and the DOM electron-donating capacity (EDC), which is mainly due to polyphenol moieties in DOM (McKay et al., 2017). This lack of good correlation could be due to the small experimental database, the heterogeneity in producing HO· from different substituted phenols (Sun et al., 2015), and the fact that some polyphenolic HO· precursors are formed only after DOM is irradiated (Berto et al., 2016). Thus, further studies are needed to clarify the production mechanism of HO· in DOM solutions and, in

particular, the correlation between the quantum yield for HO \cdot production and EDC should be investigated using irradiated DOM.

4.2.2 Inhibition effects of DOM

Light screening and quenching are the two main inhibition effects of DOM on the photodegradation of OMPs. The light screening effect is based on the fact that DOM competes for light with the OMP resulting in inhibition of the direct photolysis. The light screening effect is unavoidable for most OMPs because direct photolysis is induced mainly by short-wavelength light, which overlaps the absorption spectra of DOM under solar radiation. In addition to OMPs, DOM also screens and weakens radiation for other photosensitizers such as NO $_3^-$ /NO $_2^-$, inhibiting their photolysis that yields HO \cdot .

Phenolic moieties are mainly responsible for the quenching effects in DOM solutions. These electron-rich compounds can transfer an electron to $^3\text{DOM}^*$, thereby consuming it (Canonica et al., 2000), but this inhibition effect is not predominant in sunlit natural water bodies (Canonica et al., 2000). A recent study reported that quenching of Hi- $^3\text{DOM}^*$ is more apparent than that of low-energy $^3\text{DOM}^*$ owing to the higher redox potential of Hi- $^3\text{DOM}^*$ (Zhou et al., 2019a). On the other hand, the phenolic moieties can reduce the reactive intermediates generated from the oxidation reaction between $^3\text{DOM}^*$ and OMPs. For example, oxidation of aromatic anilines by $^3\text{DOM}^*$ forms a cation radical (aromatic aniline after loss of an electron). Electron-rich phenolic moieties can transfer an electron to the cation radical reducing it back to its parent compound, thus inhibiting the reaction (Wenk and Canonica, 2012). Previous studies have reported that the content of phenolic moieties in autochthonous DOM is lower than that in allochthonous DOM (Guo et al., 2021; Zhou et al., 2019a), and thus the quenching effects is less obvious in autochthonous DOM (Bahnmüller et

al., 2014; Wang et al., 2019; Wenk and Canonica, 2012). Additionally, Canonica and Laubscher (2008) reported that quenching effects were observed in the photodegradation of OMPs containing phenols, anilines, and on the like and that the inhibition efficiencies were different for different compounds.

Photobleaching of DOM can be considered an indirect inhibition effect of DOM as well. After photobleaching of DOM, its photo-reactivity is decreased (Helms et al., 2009, 2014; Moran et al., 2000; Niu et al., 2014). In photobleaching, the chromophores are photochemically destroyed either through direct destruction of the DOM molecules by light or reactions between DOM and PPRI. After photobleaching of DOM, the rate of photodegradation of OMPs in the same solutions could decrease. Niu et al. (2014) reported that after 20 h of simulated sunlight exposure, the observed photodegradation rate constant of some OMPs decreased as the length of irradiation time increased, and the decrease was highly dependent on the characteristics of both the DOM and OMPs. Using solid-state ^{13}C nuclear magnetic resonance spectroscopy and Fourier transform infrared spectroscopy, Helms et al. (2014) found that aromatic carbon was preferentially removed while carbohydrate-like and amide/peptide-like carbons remained after photobleaching for up to 110 days.

4.2.3 Seawater DOM vs freshwater DOM

Although seawater covers 71% of the Earth's surface and acts as an important sink of OMPs, less attention has been paid to the effects of seawater DOM (S-DOM) on the photodegradation of OMPs compared with freshwater DOM. This is due to the difficulty of extracting S-DOM from seawater, which typically has high salinity and a relatively low DOM content. S-DOM is mainly autochthonous DOM that originates from cellular excretions of marine phytoplankton and bacteria; S-DOM normally undergoes photobleaching for a long time and can be influenced by anthropogenic

sources such as mariculture. Hence, the effects of S-DOM on photodegradation of OMPs can differ from those of freshwater DOM.

Reverse osmosis coupled with electro dialysis (RO/ED) technology is normally used to extract S-DOM because the efficiency of recovery is high (Zhang et al., 2013). Wang et al. (2018) extracted S-DOM by RO/ED and found that S-DOM influenced by mariculture activities exhibited higher photo-reactivity (higher PPRIs production quantum yield, light absorption rate and $^3\text{DOM}^*$ reactivity) than S-DOM extracted from more pristine seawater. Wang et al. (2019) also found that compared with reference freshwater DOM (Suwannee River Fulvic Acid and Suwannee River Natural Organic Matter (SRNOM)), S-DOM undergoes more photobleaching in which a lower content of C=C and C=O functional groups is observed. However, compared with freshwater DOM, S-DOM enhanced the photodegradation of both 2,4-dihydroxybenzophenone and nebulolol to a greater extent, which was attributed to the higher reactivity of $^3\text{DOM}^*$ with the OMP and fewer antioxidants in the S-DOM (Wang et al., 2020b; Wang et al., 2019). Further work by Guo et al. (2021) indicated that $\text{Hi-}^3\text{DOM}^*$ is not responsible for the higher reactivity of $^3\text{DOM}^*$ from S-DOM than that from freshwater DOM. In contrast, a lower content of phenolic moieties that lessened inhibition explained the higher reactivity of $^3\text{DOM}^*$ from S-DOM compared with freshwater DOM. However, the S-DOM was extracted only from coastal areas in the Bohai Sea and further studies concerning the photochemistry of S-DOM from other regions are needed.

4.3 Suspended solids and coexisting pollutants

Suspended solids such as sand, clay, particulate organic matter, and phytoplankton are also ubiquitous in surface waters. These components can influence the underwater light environment, adsorb OMPs, and affect photosensitizing

processes (Kirk, 1985; Lund-Hansen, 2004; Rose et al., 2014; Shang et al., 2015), possibly altering the photodegradation of OMPs. For example, Niu et al. (2013) found that suspended solids obtained from a river inhibited the photodegradation of sulfamethoxazole and the inhibition effects became stronger with increasing concentration of the suspended sediments. The inhibition effect was mainly due to the suppression of HO \cdot generation in the suspended sediments solutions while adsorption of sulfamethoxazole on solids played an insignificant role. Thirumavalavan et al. (2012) attributed the effect of suspended solids to turbidity and light attenuation, and their study showed that the photodegradation of microcystin-LR was impeded when the turbidity increased.

Notably, a variety of OMPs, rather than a single one, may coexist in the same water body and thus the photodegradation of an OMP can be influenced by the coexisting OMPs. Generally, the coexisting OMPs can have multifarious effects on the photodegradation processes (Jentzsch et al., 2019; Zhang et al., 2018b) depending on the intended functions of the coexisting OMPs. For example, UV absorbers (used in polymers) are designed to absorb photon energy and then to release the excitation energy in the form of heat, so they can compete for photons with the target OMP during photodegradation. Primary antioxidants are capable of reacting with PPRIs, thus preventing further propagation of oxidation reactions. The antioxidants can also consume the generated PPRIs and prevent further reactions with the target OMP (Hunt, 2000).

In addition, recent studies have reported that trace amounts of co-existing OMPs can also increase the photodegradation of other OMPs in waters (Jentzsch et al., 2019; Zhang et al., 2018b; Zhou et al., 2019b). For example, bezafibrate (BZF) displayed faster photodegradation in the presence of fenofibrate acid (FNFA) in ultrapure water,

wastewater treatment plant effluent water, and the receiving river water, despite the usual resistance of BZF to degradation. The promotional effect of FNFA was due to its ability to generate $^1\text{O}_2$ and e^-_{aq} . Further, tests conducted with SRNOM (5.0 and 10.0 mg/L) and BZF (5.0 mg/L) in the presence and absence of FNFA (1.0 mg/L) revealed that the presence of FNFA facilitated much faster photodegradation of BZF than the presence of SRNOM (Zhou et al., 2019b).

Hence, the influence of co-existing OMPs on each other's degradation can be important in surface waters, despite their lower concentration compared with DOM. However, the influences of co-exposure of the OMPs on each other's photodegradation has hardly been studied in water environments. Given the increased detection of diverse OMPs in surface waters, further investigations are necessary to elucidate the influence of co-existing OMPs on each other's persistence in environmental waters.

5. Major physical factors for photodegradation of OMPs

Photodegradation is also influenced by physical environmental factors, such as the depth of water, latitude, temporal variation (diurnal and seasonal) and temperature (Koumaki et al., 2015; McKay and Rosario-Ortiz, 2015; Zepp and Cline, 1977). Those physical effects have received less attention than the water matrix because the physical effects are basically caused by variation of the light intensity and the wavelength profile apart from the diversity of photochemical reactions. However, changes in physical factors cannot be ignored when aiming to understand the photochemical reactions and the overall fate of OMPs. For example, thermal lake stratification can determine the vertical distribution of OMP concentrations as a result of the vertical mixing process and photolysis in the surface layer (Rehrl et al., 2020). Assuming the same wavelength profile, the rate of photodegradation would be

proportional to the photon influx. In fact, most studies involving *in situ* photodegradation experiments performed at different locations and different times (i.e., different latitudes and seasons), might be more readily comparable if the effects of physical factors on light conditions are better understood.

5.1 Latitude

Sunshine duration and cloud conditions vary depending on latitude, and clearly this would influence the availability of underwater light and thus light absorption by photosensitizers or OMPs. Annual irradiance (measured over the period 1951–1960) shows a substantial change with latitude (Figure 4) (Landsberg, 1961; Lewis, 1987).

Total annual irradiance on the Earth's surface is distributed around a maximum at the equator. Theoretically, the tropical latitudes (0° to 25°) should have the highest annual irradiance (Figure 4, theoretical Earth surface) because the sun shines almost directly over these areas. However, in fact, subtropical latitudes (25° to 40°) have the highest annual irradiance (Figure 4, land masses). This is because cloud conditions can influence the amount of solar radiation that reaches the Earth's surface. Clouds can decrease the solar flux by up to 70% of the initial solar radiation that is transmitted through the atmosphere to the surface (Janjai et al., 2011). The effect of cloud cover on decreasing light transmission has been estimated to be between -31% for UVB radiation and -72% for photosynthetically active radiation (PAR) (Musiolková et al., 2021). In addition, the uneven distribution of moisture causes the transmissivity of the atmosphere to be unusually high over subtropical land masses. Thus, more light can reach the Earth's surface at subtropical latitudes than in tropical regions (Lewis, 1987).

Given a constant attenuation coefficient, daily irradiance at latitudes between 10°N and 5°S shows peaks twice a year in March and September (Lewis, 1987). The

annual minimum daily irradiance for a constant attenuation coefficient near the equator would be approximately 85% of the annual maximum daily irradiance. The range of minima on the tropical latitudes is significantly higher, and the minimum daily irradiance at 30° latitude is approximately 50% of the minimum at the equator (Figure 4). The relative importance of shorter wavelengths also varies depending on the latitude, which possibly affects the absorption spectra and the rate of photodegradation of OMPs (Schmitt et al., 2021).

Due to the tilt of the Earth's axis of rotation, solar energy is received at a low angle at high latitudes; hence, the distribution of net energy per unit area is much less than that in the tropics. Also, higher latitudes (above 50°) often have the shortest sunshine duration annually and have more cloudy and rainy weather. At latitudes higher than 66.7° (the polar region), no direct energy is received at all from the sun for at least part of the year (Roots, 1989). Therefore, the rate of degradation rate and the half-life times that are determined experimentally need to be corrected to estimate the actual degradation rate in the different regions. Furthermore, the attenuation of light by clouds has uncertainties with respect to the climatic reduction of light intensity at various geographical locations.

Indirect relationships of latitude with the water matrix and photochemistry can be identified based on the environmental conditions of the surface water. For instance, lakes located in high-latitude regions may be covered with ice for much of the year, so sunlight can barely reach the underlying water. Until the ice has melted, the photochemical process may possibly cease due to the efficient attenuation of light (Sommaruga, 2001). Moreover, several surface waters in high-latitude Nordic countries experience browning, which is a gradual increase in DOM concentration (Weyhenmeyer et al., 2016). Such browning could affect photoreactions by enhancing

processes that are triggered by $^3\text{DOM}^*$ and $^1\text{O}_2$, over those induced by $\text{HO}\cdot$, CO_3^- and direct photolysis (Calderaro and Vione, 2020). In addition to shifting the photodegradation pathways toward $^3\text{DOM}^*$, the increase in DOM increase could also lower the photodegradation rate constants and increase the photochemical persistence of OMPs (Vione and Scozzaro, 2019). Recent studies have updated the solar irradiance spectra between 70°S and 70°N at different times of the year, and these were validated against high-resolution spectroradiometer measurements (Apell and McNeill, 2019). Simulations of PPRI production using the updated solar irradiance spectra show that the spectrum of sunlight plays a role that is as important as the role of light intensity in determining PPRI production. Furthermore, the dependence of PPRI production on the solar irradiance spectra has recently been evaluated by designing reactors equipped with monochromatic LED lights for the experiments. The result show that the quantum yields of RIs decreased with increasing wavelength from 375 to 490 nm and, notably, decreased to zero above 490 nm (Wu et al., 2021).

5.2 Seasonal and diurnal cycles

The solar irradiance varies daily and seasonally due to the rotation and revolution of the Earth, and the temporal variation of sunlight affects the photodegradation of OMPs. In terms of daily variation, the photochemical persistence of OMPs can be prolonged (for OMPs with photodegradation $t_{1/2}$ being over days) because no photodegradation occurs at nighttime. For example, recent studies indicated that the photodegradation $t_{1/2}$ of OMPs can be significantly underestimated if the variation of the intensity of sunlight with time of day is not considered (Zhou et al., 2018; Chen et al., 2019; Guo et al., 2021). On the other hand, the variations (i.e., seasonal and diurnal) of the water matrix such as pH, NO_2^- , NO_3^- , and DOM together with the irradiance of sunlight, all affected the photochemical processes including the overall

kinetics of the OMP photo-transformations (Vione et al., 2018). Compared with freshwater and estuarine systems, the attenuation of sunlight in seawater is normally the weakest (Zhou et al., 2018), and thus such differences will affect photochemical processes and rates.

The diurnal solar light variations in Nigeria (tropic) (Soneye et al., 2019) and Northeast China (mid-latitude) (Zhou et al., 2018) are presented in Figure 5A and 5B, respectively. The solar intensity increases from 7 am, reaching a maximum at around noon, and decreases to 0 Wm^{-2} at around 6 pm. The variation in intensity of solar light suggested that the photodegradation rate of OMPs should likewise show a variation, although photodegradation studies usually simulate a solar intensity at noon which could largely overestimate the degradation rate of the OMPs. In addition, as shown in Figure 5A, variation of the diurnal pattern of light intensity over the seasons can also be observed. Such differences in light intensity are caused by the attenuation of light by aerosol particles in the dry season and increased cloudiness and humidity in the wet season.

Stratification is also a seasonal phenomenon that commonly occurs in lakes and reservoirs, which also influences the photochemical processes in waters. For example, lake water typically undergoes stratification in summer in temperate climates. The epilimnion is the surface layer of water and thus receives more illumination, meaning that photodegradation processes will be more active there. In contrast, the hypolimnion is poorly illuminated due to light screening by the overlying water column, so photochemical processes are limited or negligible (Vione, 2020). The epilimnion and hypolimnion evolve differently, and thus a depth profile gradient is generated with respect to the concentration of photodegradable OMPs (Avetta et al., 2016). If the OMP is completely photodegraded in the epilimnion, the stratification

phase considerably slows down the overall degradation kinetics (Vione and Scozzaro, 2019).

A recent study investigated the photodegradation of 23 OMPs in the free-water surface of a constructed wetland basin during the summer and winter in southern France (43.85°N). For the 23 OMPs studied, direct photolysis was the predominant process in summer, while indirect photodegradation was more significant in winter (Mathon et al., 2019). Although the light intensity was higher in summer, the NO_3^- concentration was higher in winter enhancing the formation of HO which promoted the wintertime indirect photodegradation (Mathon et al., 2019). In addition, a 7-year survey in a stream in the USA observed seasonal variation of NO_3^- in which high NO_3^- concentrations were observed in summer and low NO_3^- concentrations were observed from mid-autumn to late spring. The low NO_3^- concentrations were attributed to maximum sunlight irradiance occurring in late spring and the seasonal input of leaves to the stream in autumn (Mulholland and Hill, 1997). Another study conducted in the Guadiana River, Spain, has also shown that photodegradation is important in summer, and quite slow in winter (Vione et al., 2018). In the months with the highest clear-sky sunlight irradiance (June and July), the river water showed low NO_3^- concentrations and high DOM levels. This phenomenon may be quite typical of surface water bodies as the minima for NO_3^- and the maxima for DOC occur frequently in summer in several surface-water environments (Vione et al., 2018).

5.3 Water Depth

Underwater light intensity is generally attenuated with increasing water depth due to components such as phytoplankton, particles, DOM, ions, and water itself. H_2O absorbs mainly red light (around 700 nm), which is one of the reasons for the color of

the sea (Hale and Querry, 1973). Sunlight attenuation underwater alters the availability of light and affects the photochemical reactions of OMPs and photosensitizers (Zhou et al., 2018). Light absorption and scattering are the main causes of light attenuation in water bodies (Kirk, 1977). In general, the intensity of light decreases exponentially with depth as described by

$$I_z = I_0 e^{-Ez} \quad (14)$$

where I_z and I_0 are the intensities of light (W m^{-2} or quanta $\text{m}^{-2} \text{s}^{-1}$) at depth z (m) and just below the surface, respectively, and E is the vertical attenuation coefficient (m^{-1}) for downward irradiance (Scheffer, 2004), which is affected by water matrix components such as DOM and inorganic ions. The depth of this photic environment varies depending on the water matrix and the wavelength of light (Avetta et al., 2016; Boreen et al., 2003; Vione and Scozzaro, 2019). The depth of light penetration generally refers to the depth at which the light has been attenuated by 99% (Zepp and Cline, 1977). Turbid lakes and estuaries show a light penetration depth of < 0.3 m, while the clearest oceans have a penetration depth up to 150 m (Gons et al., 1997). In terms of wavelength, the depth of light penetration in freshwater is low for UV, blue, and red light, whereas green and yellow transmit to deeper water (Levine and MacNichol, 1982).

Moreover, the wavelength profile of irradiation shifts vertically in water depending on the water matrix (Minero et al., 2007). The absorption effects of DOM change depending on the wavelength profile, especially for light with wavelengths below 500 nm. Therefore, penetration depth for sunlight is greater in waters containing lower concentrations of DOM and the corresponding rates of photo-transformation may be larger (Bracchini et al., 2007). In addition, the enhancement of

photoreactions in shallow water is more conspicuous for processes triggered by UV absorbing species such as $\text{NO}_3^-/\text{NO}_2^-$ (Vione and Scozzaro, 2019).

Previous studies concluded that UV light is attenuated by water within tens of centimeters of depth in lakes with a high DOM concentration and in turbid waters; in both cases the photochemical reactions are significantly affected by the light attenuation (Huovinen et al., 2003; Leavitt et al., 1997; Markager and Vincent, 2000). Another study reported that the steady-state concentration of $^1\text{O}_2$ in lake waters decreased by a factor of 8.8 when the depth increased from 0 m to 20 m, indicating that depth is an important factor in determining indirect photochemical processes (Partanen et al., 2021). The APEX model can be used to predict variation in the rate of photodegradation with depth, and one study employing this model demonstrated that the photochemical half-lives $t_{1/2}$ of three OMPs (ibuprofen, carbamazepine, and diclofenac) are almost linearly correlated with depth (Vione et al., 2018).

In addition to the variation of light attenuation with depth, photochemical reactions in the surface layer ($z = 0$ m) differ from those in bulk water, because the surface layer (air/water) creates a unique environment that can promote unique chemical reactions (George et al., 2015; Kappes et al., 2021). For example, interfaces covered with fatty acids are ubiquitous in the environment and fatty acids are believed to be photochemically inert in the actinic region; however, direct photolysis of fatty acids has been shown to occur, and complex volatile organic compounds are generated at an air-water interface coated solely with a monolayer of carboxylic acid (Rossignol et al., 2016). Likewise, photochemical reactions may also occur at the water/sediment interface in water bodies where the light is not completely attenuated at the bottom of the water column. These photochemical water/sediment interface reactions might determine the fate of many hydrophobic OMPs because sorption by

sediments is an important removal pathway for OMPs (Golovko et al., 2020; Koba et al., 2018). Nevertheless, the photochemical reactions at interfaces have received relatively little attention and further studies are needed.

5.4 Temperature

Generally, temperature is higher in summer than in winter and decreases with increasing water depth and latitude. Several photodegradation studies have highlighted the impact of temperature on the photodegradation process. McKay and Rosario-Ortiz (2015) studied the effect of temperature on HO· formation using five different types of DOM isolates at four or five different temperatures within the range 10-40 °C. The apparent activation energies E_a for HO· formation from the DOMs (kJ mol⁻¹) were obtained using the Arrhenius equation (Eq. 15)

$$\ln(\Phi_a) = -E_a/RT + \ln(A) \quad (15)$$

where Φ_a is the apparent quantum yield of HO· through DOM photolysis, R is the universal gas constant (kJ K⁻¹ mol⁻¹), T is the absolute temperature (K), and A is the pre-exponential factor. Plots of $\ln(\Phi_a)$ vs (1/T) revealed significant slopes, indicating the temperature dependency of HO· formation through photolysis of the DOM isolates.

In addition, the photolysis of *p*-benzoquinone was studied by applying the Arrhenius equation to the production of phenol through *p*-benzoquinone photolysis and the loss of *p*-benzoquinone. It was revealed that Φ_a of formation of phenol through *p*-benzoquinone photolysis was temperature independent while Φ_a of *p*-benzoquinone loss was temperature independent when the temperature was over 23 °C and temperature dependent when the temperature was less than 23 °C (McKay and Rosario-Ortiz, 2015).

Zhu and Kieber (2018) studied the photolysis of acetaldehyde, glyoxal, and methylglyoxal in North Pacific seawater under simulated solar radiation and revealed that the apparent quantum yield of acetaldehyde was temperature independent while those for glyoxal and methylglyoxal were temperature dependent. Another study was conducted by Kieber et al. (2014) to determine the impact of wavelength and temperature on the apparent quantum yield for producing H_2O_2 with seawater, photochemically. Seawater from stations located in Antarctica, the Pacific Ocean, the Gulf of Mexico and at a number of sites located on the East Coast of the United States were examined. The experiments revealed that the apparent quantum yields for H_2O_2 production increased within the temperature range 0–35⁰C with an average factor of 1.8 every 10⁰C. The study indicated that the decrease of temperature and irradiation toward the polar region was mainly responsible for the decrease in H_2O_2 production in the seawater.

As indicated above, temperature affects the photochemical transformations in the aquatic environment even though the influence varies depending on the OMP or the process of concern. Therefore, it is necessary to acknowledge temperature as an important physical factor influencing photochemical reactions in water.

6. Major knowledge gaps and challenges

- Studies of underwater light attenuation have focused mainly on PAR (i.e., 400–700 nm) while less attention has been paid to the UV region (i.e., 280–400 nm), which is critically important for photodegradation. This bias hinders better understanding of photochemical processes in aquatic environments.
- The DOM photochemistry is complex and has not been comprehensively explored. Further investigations are needed with a specific focus on topics such

as the production mechanisms of $\text{HO}\cdot$, the role of $\text{H}^+\text{-}^3\text{DOM}^*$ in photodegradation, and the photo-reactivity of less-studied DOM (e.g., seawater DOM).

- Modeling of photodegradation rates of OMPs in various water bodies is required for comprehensive understanding and proper management of OMPs in water environments. Although the APEX model is kinetics-based and practical, there is scope for further development, such as comprehensive studies on photodegradation in actual water systems, for instance with consideration of physical factors and the reactivity of real $^3\text{DOM}^*$.
- Further research efforts are needed to develop prediction models for parameters necessary to describe the kinetics of OMP photodegradation (quantum yield, second-order rate constant, etc.), so as to improve model applicability in terms of domain and prediction accuracy (at present it is impossible to exhaustively determine the kinetics parameters of all OMPs experimentally).
- Photodegradation has been shown to enhance the toxicity of OMPs in some cases, so researchers must also pay attention to degradation pathways and the potential toxicity of byproducts (Vione and Carena, 2020). Computational toxicology can be of great help in assessing the ecotoxicity of products as standard chemicals are not always available.
- The available photochemistry studies are biased toward temperate and subarctic regions, and thus more attention should be paid to tropical regions where the incident solar radiation is more intense and many developing countries may have to deal with serious pollution caused by OMPs.
- A standard protocol for indirect photodegradation studies is needed to provide meaningful data to allow comparative research and ecological risk assessment.

Recommendations for the determination of $^1\text{O}_2$ quantum yield is one such example (Ossola et al., 2021).

- In general, the increasing diversity of OMPs in water bodies is responsible for multifarious photodegradation, which poses challenges both in environmental science and environmental management for maintaining sustainable aquatic ecosystems.

7. Conclusions

Chemical pollution by OMPs has become a serious environmental issue and understanding the fate of OMPs is essential for conducting ecological risk assessments. This review has highlighted the fact that photodegradation is a major removal process for many OMPs in the aquatic environment, and its significance and processes depend on chemical and physical conditions. The major factors determining these conditions include the water matrix such as inorganic ions, DOM, suspended solids, and coexisting pollutants, and physical factors such as latitude, depth, temporal variation of light intensity, and temperature. Much attention has been given to the complex effects of DOM and the recent advancements in DOM photochemistry. Finally, we identified major knowledge gaps and challenges, which we hope will further stimulate further investigation of photodegradation processes and the fate of OMPs and, in so doing, bring about improved understanding and control of OMPs in environmental waters.

Acknowledgment

This review is one of the outcomes from projects funded by MEXT/JSPS (KAKENHI, 21H01462) and the Water Resources Environmental Center. Zhongyu Guo

acknowledges financial support from the Advanced Human Resource Development Fellowship for Doctoral Students, Tokyo Institute of Technology.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- Arakaki, T., Miyake, T., Hirakawa, T., Sakugawa, H. 1999. pH dependent photoformation of hydroxyl radical and absorbance of aqueous-Phase N(III) (HNO_2 and NO_2^-). *Environ. Sci. Technol.* 33, 2561-2565. <https://doi.org/10.1021/es980762i>
- Avetta, P., Fabbri, D., Minella, M., Brigante, M., Maurino, V., Minero, C., Pazzi, M., Vione, D., 2016. Assessing the phototransformation of diclofenac, clofibric acid and naproxen in surface waters: Model predictions and comparison with field data. *Water Res.* 105, 383–394. <https://doi.org/10.1016/j.watres.2016.08.058>
- Baena-Nogueras, R.M., González-Mazo, E., Lara-Martín, P.A., 2017. Degradation kinetics of pharmaceuticals and personal care products in surface waters: photolysis vs biodegradation. *Sci. Total Environ.* 590–591, 643–654. <https://doi.org/10.1016/j.scitotenv.2017.03.015>
- Bahnmüller, S., von Gunten, U., Canonica, S., 2014. Sunlight-induced transformation of sulfadiazine and sulfamethoxazole in surface waters and wastewater effluents. *Water Res.* 57, 183–192. <https://doi.org/10.1016/j.watres.2014.03.019>
- Benedict, K.B., McFall, A.S., and Anastasio, C., 2017. Quantum yield of nitrite from the photolysis of aqueous nitrate above 300 nm. *Environ. Sci. Technol.* 51, 4387-4395. <https://doi.org/10.1021/acs.est.6b06370>

- Berto, S., Laurentiis, E.D., Tota, T., Chiavazza, E., Daniele, P.G., Minella, M., Isaia, M., Brigante, M., Vione, D., 2016. Properties of the humic-like material arising from the photo-transformation of L-tyrosine. *Sci. Total. Environ.* 545-546, 434-444. <https://doi.org/10.1016/j.scitotenv.2015.12.047>
- Bodrato, M., Vione, D., 2014. APEX (Aqueous Photochemistry of Environmentally occurring Xenobiotics): A free software tool to predict the kinetics of photochemical processes in surface waters. *Environ. Sci. Process. Impacts* 16, 732–740. <https://doi.org/10.1039/c3em00541k>
- Boreen, A.L., Arnold, W.A., McNeill, K., 2003. Photodegradation of pharmaceuticals in the aquatic environment: A review. *Aquat. Sci.* 65, 320–341. <https://doi.org/10.1007/s00027-003-0672-7>
- Boreen, A.L., Arnold, W.A., McNeill, K., 2004. Photochemical fate of sulfa drugs in the aquatic environment: sulfa drugs containing five-membered heterocyclic groups. 38, 3933-3940. <https://doi.org/10.1021/es0353053>
- Boreen, A.L., Edhlund, B.L., Cotner, J.B., McNeill, K., 2008. Indirect photodegradation of dissolved free amino acids: The contribution of singlet oxygen and the differential reactivity of DOM from various sources. *Environ. Sci. Technol.* 42, 5492–5498. <https://doi.org/10.1021/es800185d>
- Bouhala, A., Lahmar, H., Benamira, M., Moussi, A., Trari, M., 2020. Photodegradation of organophosphorus pesticides in honey medium by solar light irradiation. *Bull. Environ. Contam. Toxicol.* 104, 792–798. <https://doi.org/10.1007/s00128-020-02858-1>
- Boule, P. 1999. *Environmental Photochemistry, The Handbook of Environmental Chemistry*. Springer, Berlin, Heidelberg. <https://doi.org/10.1007/978-3-540-69044-3>
- Boyle, E.S., Guerriero, N., Thiallet, A., Del Vecchio, R., Blough, N. V., 2009. Optical properties of humic substances and CDOM: Relation to structure. *Environ. Sci. Technol.* 43, 2262-2268. <https://doi.org/10.1021/es803264g>

- Bracchini, L., Loisel, S., Dattilo, A.M., Mazzuoli, S., C zar, A., Rossi, C., 2007. The Spatial Distribution of Optical Properties in the Ultraviolet and Visible in an Aquatic Ecosystem. *Photochem. Photobiol.* 80, 139–149. <https://doi.org/10.1111/j.1751-1097.2004.tb00062.x>
- Brausch, J.M., Rand, G.M., 2011. A review of personal care products in the aquatic environment: Environmental concentrations and toxicity. *Chemosphere* 82, 1518–1532. <https://doi.org/10.1016/j.chemosphere.2010.11.018>
- Burns, S.E., Hassett, J.P., Rossi, M.V., 1996. Binding effects on humic-mediated photoreaction: Intrahumic dechlorination of mirex in water. *Environ. Sci. Technol.* 30, 2934–2941. <https://doi.org/10.1021/es950906i>
- Burns, S.E., Hassett, J.P., Rossi, M.V., 1997. Mechanistic implications of the intrahumic dechlorination of mirex. *Environ. Sci. Technol.* 31, 1365–1371. <https://doi.org/10.1021/es960581i>
- Busset, C., Mazellier, P., Sarakha, M., De Laat, J., 2007. Photochemical generation of carbonate radicals and their reactivity with phenol. *J. Photochem. Photobiol. A Chem.* 185, 127–132. <https://doi.org/10.1016/j.jphotochem.2006.04.045>
- Buth, J.M., Arnold, W.A., McNeill, K., 2007. Unexpected products and reaction mechanisms of the aqueous chlorination of cimetidine. *Environ. Sci. Technol.* 41, 6228–6233. <https://doi.org/10.1021/es070606o>
- Calderaro, F., Vione, D., 2020. Possible effect of climate change on surface-water photochemistry: A model assessment of the impact of browning on the photodegradation of pollutants in lakes during summer stratification. Epilimnion vs. whole-lake phototransformation. *Molecules* 25, 1–14. <https://doi.org/10.3390/molecules25122795>
- Calza, P. and Vione, D. 2015. Surface Water Photochemistry, Comprehensive Series in Photochemical & Photobiological Sciences. Royal Society of Chemistry, Cambridge, UK. <https://doi.org/10.1039/9781782622154>

- Canonica, S., Hellrung, B., Wirz, J., 2000. Oxidation of phenols by triplet aromatic ketones in aqueous solution. *J. Phys. Chem. A* 104, 1226–1232. <https://doi.org/10.1021/jp9930550>
- Canonica, S., Kohn, T., Mac, M., Real, F.J., Wirz, J., Von Gunten, U., 2005. Photosensitizer method to determine rate constants for the reaction of carbonate radical with organic compounds. *Environ. Sci. Technol.* 39, 9182–9188. <https://doi.org/10.1021/es051236b>
- Canonica, S., Laubscher, H.U., 2008. Inhibitory effect of dissolved organic matter on triplet-induced oxidation of aquatic contaminants. *Photochem. Photobiol. Sci.* 7, 547–551. <https://doi.org/10.1039/b719982a>
- Challis, J.K., Carlson, J.C., Friesen, K.J., Hanson, M.L., Wong, C.S., 2013. Aquatic photochemistry of the sulfonamide antibiotic sulfapyridine. *J. Photochem. Photobiol. A Chem.* 262, 14–21. <https://doi.org/10.1016/j.jphotochem.2013.04.009>
- Challis, J.K., Hanson, M.L., Friesen, K.J., Wong, C.S., 2014. A critical assessment of the photodegradation of pharmaceuticals in aquatic environments: Defining our current understanding and identifying knowledge gaps. *Environ. Sci. Process. Impacts* 16, 672–696. <https://doi.org/10.1039/c3em00615h>
- Chen, J.W., Peijnenburg, W.J.G.M., Quan, X., Yang, F.L., 2000. Quantitative structure-property relationships for direct photolysis quantum yields of selected polycyclic aromatic hydrocarbons. *Sci. Total. Environ.* 246, 11–20. [https://doi.org/10.1016/S0048-9697\(99\)00407-6](https://doi.org/10.1016/S0048-9697(99)00407-6)
- Chen, J.W., Peijnenburg, W.J.G.M., Quan, X., Chen, S., Martens, D., Schramm, K.-W., Kettrup, A., 2001a. Is it possible to develop a QSPR model for direct photolysis half-lives of PAHs under irradiation of sunlight? *Environ. Pollut.* 114, 137–143. [https://doi.org/10.1016/S0269-7491\(00\)00195-0](https://doi.org/10.1016/S0269-7491(00)00195-0)
- Chen, J.W., Quan, X., Peijnenburg, W.J.G.M., Yang, F.L., 2001b. Quantitative structure–property relationships (QSPRs) on direct photolysis quantum yields of

- PCDDs. *Chemosphere* 43, 235-241. [https://doi.org/10.1016/S0045-6535\(00\)00141-7](https://doi.org/10.1016/S0045-6535(00)00141-7)
- Chen, X., Wang, J.Q., Chen, J.W., Zhou, C.Z., Cui, F.F., Sun, G.X., 2019. Photodegradation of 2-(2-hydroxy-5-methylphenyl) benzotriazole (UV-P) in coastal seawaters: Important role of DOM. *J. Environ. Sci. (China)* 85, 129-137. <https://doi.org/10.1016/j.jes.2019.05.017>
- Cheng, F.Y., He, J.L., Li, C., Lu, Y., Zhang, Y.N., Qu, J., 2021. Photo-induced degradation and toxicity change of decabromobiphenyl ethers (BDE-209) in water: Effects of dissolved organic matter and halide ions. *J. Hazard. Mater.* 416, 125842. <https://doi.org/10.1016/j.jhazmat.2021.125842>
- Coddington, O., Lean, J.L., Pilewskie, P., Snow, M., Lindholm, D., 2016. A solar irradiance climate data record. *Bull. Am. Meteorol. Soc.* 97, 1265–1282. <https://doi.org/10.1175/BAMS-D-14-00265.1>
- Cooper, W.J., Lean, D.R., 1989. Hydrogen peroxide concentration in a northern lake: photochemical formation and diel variability. *Environ. Sci. Technol.* 23, 1425-1428. <https://doi.org/10.1021/es00069a017>
- Crane, M., Watts, C., Boucard, T., 2006. Chronic aquatic environmental risks from exposure to human pharmaceuticals. *Sci. Total Environ.* 367, 23–41. <https://doi.org/10.1016/j.scitotenv.2006.04.010>
- De la Cruz, N., Esquius, L., Grandjean, D., Magnet, A., Tungler, A., de Alencastro, L.F., Pulgarín, C., 2013. Degradation of emergent contaminants by UV, UV/H₂O₂ and neutral photo-Fenton at pilot scale in a domestic wastewater treatment plant. *Water Res.* 47, 5836–5845. <https://doi.org/10.1016/j.watres.2013.07.005>
- De Laurentiis, E., Prasse, C., Ternes, T.A., Minella, M., Maurino, V., Minero, C., Sarakha, M., Brigante, M., Vione, D., 2014. Assessing the photochemical transformation pathways of acetaminophen relevant to surface waters:

- Transformation kinetics, intermediates, and modelling. *Water Res.* 53, 235–248.
<https://doi.org/10.1016/j.watres.2014.01.016>
- Dimou, A.D., Sakkas, V.A., Albanis, T.A., 2005. Metolachlor photodegradation study in aqueous media under natural and simulated solar irradiation. *J. Agric. Food Chem.* 53, 694–701. <https://doi.org/10.1021/jf048766w>
- Ebert, I., Bachmann, J., Kühnen, U., Küster, A., Kussatz, C., Maletzki, D., Schlüter, C., 2011. Toxicity of the fluoroquinolone antibiotics enrofloxacin and ciprofloxacin to photoautotrophic aquatic organisms. *Environ. Toxicol. Chem.* 30, 2786–2792. <https://doi.org/10.1002/etc.678>
- Edlund, B.L., Arnold, W.A., McNeill, K., 2006. Aquatic photochemistry of nitrofurantoin antibiotics. *Environ. Sci. Technol.* 40, 5422–5427.
<https://doi.org/10.1021/es0606778>
- Fatta-Kassinos, D., Vasquez, M.I., Kümmerer, K., 2011. Transformation products of pharmaceuticals in surface waters and wastewater formed during photolysis and advanced oxidation processes-Degradation, elucidation of byproducts and assessment of their biological potency. *Chemosphere* 85, 693–709.
<https://doi.org/10.1016/j.chemosphere.2011.06.082>
- Feng, W., Nansheng, D., 2000. Photochemistry of hydrolytic iron (III) species and photoinduced degradation of organic compounds. A minireview. *Chemosphere* 41, 1137–1147. [https://doi.org/10.1016/S0045-6535\(00\)00024-2](https://doi.org/10.1016/S0045-6535(00)00024-2)
- Fent, K., Weston, A.A., Caminada, D., 2006. Ecotoxicology of human pharmaceuticals. *Aquat. Toxicol.* 76, 122–159.
<https://doi.org/10.1016/j.aquatox.2005.09.009>
- Gan, D., Jia, M., Vaughan, P.P., Falvey, D.E., Blough, N. V., 2008. Aqueous photochemistry of methyl-benzoquinone. *J. Phys. Chem. A* 112, 2803–2812.
<https://doi.org/10.1021/jp710724e>
- Ge, L.K., Chen, J.W., Qiao, X.L., Lin, J., Cai, X.Y., 2009. Light-source-dependent effects of main water constituents on photodegradation of phenicol antibiotics:

- mechanism and kinetics. *Environ. Sci. Technol.* 43, 3101–3107.
<https://doi.org/10.1021/es8031727>
- Ge, L.K., Chen, J.W., Wei, X.X., Zhang, S.Y., Qiao, X.L., Cai, X.Y., Xie, Q., 2010a. Aquatic photochemistry of fluoroquinolone antibiotics: kinetics, pathways, and multivariate effects of main water constituents. *Environ. Sci. Technol.* 44, 2400–2405. <https://doi.org/10.1021/es902852v>
- Ge, L.K., Chen, J.W., Zhang, S.Y., Cai, X.Y., Wang, Z., Wang, C.L., 2010b. Photodegradation of fluoroquinolone antibiotic gatifloxacin in aqueous solutions. *Chinese Sci. Bull.* 55, 1495–1500. <https://doi.org/10.1007/s11434-010-3139-y>
- Ge, L.K., Zhang, S.Y., Xie, Q., Chen, J.W. 2010c. Progress in studies on aqueous environmental photochemical behavior of antibiotics. *Sci. Sin. Chim.* 40, 124–135. <https://doi.org/10.1360/zb2010-40-2-124>
- George, C., Ammann, M., D'Anna, B., Donaldson, D.J., Nizkorodov, S.A., 2015. Heterogeneous photochemistry in the atmosphere. *Chem. Rev.* 115, 4218–4258. <https://doi.org/10.1021/cr500648z>
- Gligorovski, S., Streckowski, R., Barbati, S., Vione, D., 2015. Environmental implications of hydroxyl radicals ($\cdot\text{OH}$). *Chem. Rev.* 115, 13051–13092. <https://doi.org/10.1021/cr500310b>
- Glover, C.M., Rosario-Ortiz, F.L., 2013. Impact of halides on the photoproduction of reactive intermediates from organic matter. *Environ. Sci. Technol.* 47, 13949–13956. <https://doi.org/10.1021/es4026886>
- Golovko, O., Rehr, A.L., Köhler, S., Ahrens, L., 2020. Organic micropollutants in water and sediment from Lake Mälaren, Sweden. *Chemosphere* 258. <https://doi.org/10.1016/j.chemosphere.2020.127293>
- Gons, H.J., Ebert, J., Kromkamp, J., 1997. Optical teledetection of the vertical attenuation coefficient for downward quantum irradiance of photosynthetically available radiation in turbid inland waters. *Aquat. Ecol.* 31, 299–311. <https://doi.org/10.1023/A:1009902627476>

- Grandbois, M., Latch, D.E., McNeill, K., 2008. Microheterogeneous concentrations of singlet oxygen in natural organic matter isolate solutions. *Environ. Sci. Technol.* 42, 9184–9190. <https://doi.org/10.1021/es8017094>
- Grebel, J.E., Pignatello, J.J., Mitch, W.A., 2011. Sorbic acid as a quantitative probe for the formation, scavenging and steady-state concentrations of the triplet-excited state of organic compounds. *Water Res.* 45, 6535–6544. <https://doi.org/10.1016/j.watres.2011.09.048>
- Grebel, J.E., Pignatello, J.J., Mitch, W.A., 2012. Impact of halide ions on natural organic matter-sensitized photolysis of 17 β -Estradiol in saline waters. *Environ. Sci. Technol.* 46, 7128-7134. <https://doi.org/10.1021/es3013613>
- Grebel, J.E., Pignatello, J.J., Song, W.H., Cooper, W.J., Mitch, W.A., 2009. Impact of halides on the photobleaching of dissolved organic matter. *Mar. Chem.* 115, 134-144. <https://doi.org/10.1016/j.marchem.2009.07.009>
- Grenni, P., Ancona, V., Barra, A., 2018. Ecological effects of antibiotics on natural ecosystems : A review. *Microchemical J.* 136, 25–39. <https://doi.org/10.1016/j.microc.2017.02.006>
- Gruchlik, Y., Linge, K., Joll, C., 2018. Removal of organic micropollutants in waste stabilisation ponds: A review. *J. Environ. Manage.* 206, 202–214. <https://doi.org/10.1016/j.jenvman.2017.10.020>
- Guo, Z.Y., Wang, J.Q., Chen, X., Cui, F.F., Wang, T.T., Zhou, C.Z., Song, G.B., Zhang, S.Y., Chen, J.W., 2021. Photochemistry of dissolved organic matter extracted from coastal seawater: Excited triplet-states and contents of phenolic moieties. *Water Res.* 188, 116568. <https://doi.org/10.1016/j.watres.2020.116568>
- Hale, G. M., and Querry, M. R. 1973. Optical constants of water in the 200-nm to 200- μ m wavelength region. *Appl. Optics* 12, 555-563. <https://doi.org/10.1364/AO.12.000555>
- Helms J.R., Stubbins, A., Ritchie, J.D., Minor, E.C., Kieber, D.J., Mopper, K., 2009. Absorption spectral slopes and slope ratios as indicators of molecular weight,

- source, and photobleaching of chromophoric dissolved organic matter. *Limnol. Oceanogr.* 54, 1023. <https://doi.org/10.4319/lo.2008.53.3.0955>
- Helms, J.R., Mao, J., Stubbins, A., Schmidt-Rohr, K., Spencer, R.G.M., Hernes, P.J., Mopper, K., 2014. Loss of optical and molecular indicators of terrigenous dissolved organic matter during long-term photobleaching. *Aquat. Sci.* 76, 353–373. <https://doi.org/10.1007/s00027-014-0340-0>
- Hernando, M.D., Mezcuca, M., Fernández-Alba, A.R., Barceló, D., 2006. Environmental risk assessment of pharmaceutical residues in wastewater effluents, surface waters and sediments. *Talanta* 69, 334–342. <https://doi.org/10.1016/j.talanta.2005.09.037>
- Huie, R.E., Clifton, C.L., Neta, P., 1991. Electron transfer reaction rates and equilibria of the carbonate and sulfate radical anions. *Int. J. Radiat. Appl. Instrument. Part.*, 38, 477–481. [https://doi.org/10.1016/1359-0197\(91\)90065-A](https://doi.org/10.1016/1359-0197(91)90065-A)
- Hunt, T.P., 2000. Polymer additives: Supercritical fluid chromatography. Level III: Practical Applications in *Encyclopedia of Separation Science*. Academic Press, UK, 3901–3906. <https://doi.org/10.1016/B0-12-226770-2/03351-2>
- Huovinen, P.S., Penttilä, H., Soimasuo, M.R., 2003. Spectral attenuation of solar ultraviolet radiation in humic lakes in Central Finland. *Chemosphere* 51, 205–214. [https://doi.org/10.1016/S0045-6535\(02\)00634-3](https://doi.org/10.1016/S0045-6535(02)00634-3)
- Janjai, S., Pankaew, P., Laksanaboonsong, J., Kitichantaropas, P., 2011. Estimation of solar radiation over Cambodia from long-term satellite data. *Renew. Energy* 36, 1214–1220. <https://doi.org/10.1016/j.renene.2010.09.023>
- Jentsch, F., Reich, M., Kümmerer, K., Olsson, O., 2019. Photolysis of mixtures of UV filters octocrylene and ethylhexyl methoxycinnamate leads to formation of mixed transformation products and different kinetics. *Sci. Total Environ.* 697, 134048. <https://doi.org/10.1016/j.scitotenv.2019.134048>
- Kameda, Y., Kimura, K., Miyazaki, M., 2011. Occurrence and profiles of organic sun-blocking agents in surface waters and sediments in Japanese rivers and lakes.

- Environ. Pollut. 159, 1570–1576.
<https://doi.org/10.1016/J.ENVPOL.2011.02.055>
- Kappes, K.J., Deal, A.M., Jespersen, M.F., Blair, S.L., Doussin, J.F., Cazaunau, M., Pangu, E., Hopper, B.N., Johnson, M.S., Vaida, V., 2021. Chemistry and photochemistry of pyruvic acid at the air-water interface. *J. Phys. Chem. A* 125, 1036–1049. <https://doi.org/10.1021/acs.jpca.0c09096>
- Kieber, D.J., Miller, G.W., Neale, P.J., Mopper, K., 2014. Wavelength and temperature-dependent apparent quantum yields for photochemical formation of hydrogen peroxide in seawater. *Environ. Sci. Process. Impacts* 16, 777–791. <https://doi.org/10.1039/C4EM00036F>
- Kirk, J.T.O., 1977. Attenuation of light in natural waters. *Mar. Freshw. Res.* 28, 497–508. <https://doi.org/10.1071/MF9770497>
- Kirk, J.T.O., 1985. Effects of suspensoids (turbidity) on penetration of solar radiation in aquatic ecosystems. *Hydrobiologia* 125, 195–208. <https://doi.org/10.1007/BF00045935>
- Klan, P., Wirz, J., 2009. *Photochemistry of Organic Compounds: From concepts to practice*. John Wiley & Sons, Ltd, Chichester, UK. <https://doi.org/10.1002/9781444300017>
- Knopp, G., Prasse, C., Ternes, T.A., Cornel, P., 2016. Elimination of micropollutants and transformation products from a wastewater treatment plant effluent through pilot scale ozonation followed by various activated carbon and biological filters. *Water Res.* 100, 580–592. <https://doi.org/10.1016/j.watres.2016.04.069>
- Koba, O., Grabicova, K., Cervený, D., Turek, J., Kolarova, J., Randak, T., Zlabek, V., Grabic, R., 2018. Transport of pharmaceuticals and their metabolites between water and sediments as a further potential exposure for aquatic organisms. *J. Hazard. Mater.* 342, 401–407. <https://doi.org/10.1016/j.jhazmat.2017.08.039>
- Köhler, G., Solar, S., Getoff, N., Holzwarth, A.R., Schaffner, K., 1985. Relationship between the quantum yields of electron photoejection and fluorescence of

- aromatic carboxylate anions in aqueous solution. *J. Photochem.* 28, 383–391.
[https://doi.org/10.1016/0047-2670\(85\)85040-1](https://doi.org/10.1016/0047-2670(85)85040-1)
- Koumaki, E., Mamais, D., Noutsopoulos, C., Nika, M.C., Bletsou, A.A., Thomaidis, N.S., Eftaxias, A., Stratogianni, G., 2015. Degradation of emerging contaminants from water under natural sunlight: The effect of season, pH, humic acids and nitrate and identification of photodegradation by-products. *Chemosphere* 138, 675–681. <https://doi.org/10.1016/j.chemosphere.2015.07.033>
- Koziar, J.C., Cowan, D.O., 1977. Photochemical heavy-atom effects. *Acc. Chem. Res.* 11, 334–341. <https://doi.org/10.1021/ar50129a003>
- Kumar, M., Jaiswal, S., Sodhi, K. K., Shree, P., Singh, D. K., Agrawal, P. K., Shukla, P., 2019. Antibiotics bioremediation: Perspectives on its ecotoxicity and resistance. *Environ. Int.* 124, 448–461.
<https://doi.org/10.1016/j.envint.2018.12.065>
- Lado Ribeiro, A.R., Moreira, N.F.F., Li Puma, G., Silva, A.M.T., 2019. Impact of water matrix on the removal of micropollutants by advanced oxidation technologies. *Chem. Eng. J.* 363, 155–173.
<https://doi.org/10.1016/j.cej.2019.01.080>
- Lallas, P.L., 2001. The Stockholm convention on persistent organic pollutants. *Am. J. Int. Law* 95, 692–708. <https://doi.org/10.2307/2668517>
- Landsberg, H.E., 1961. Solar radiation at the earth's surface. *Sol. Energy* 5, 95–98.
[https://doi.org/10.1016/0038-092X\(61\)90051-2](https://doi.org/10.1016/0038-092X(61)90051-2)
- Larsson, D.G.J., de Pedro, C., Paxeus, N., 2007. Effluent from drug manufactures contains extremely high levels of pharmaceuticals. *J. Hazard. Mater.* 148, 751–755. <https://doi.org/10.1016/J.JHAZMAT.2007.07.008>
- Latch, D.E., McNeill, K., 2006. Microheterogeneity of singlet oxygen distributions in irradiated humic acid solutions. *Science* 311, 1743–1747.
<https://doi.org/10.1126/science.1121636>

- Latch, D.E., Packer, J.L., Arnold, W.A., McNeill, K., 2003a. Photochemical conversion of triclosan to 2,8-dichlorodibenzo-*p*-dioxin in aqueous solution. *J. Photochem. Photobiol. A: Chem.* 158, 63-66. [https://doi.org/10.1016/S1010-6030\(03\)00103-5](https://doi.org/10.1016/S1010-6030(03)00103-5)
- Latch, D.E., Stender, B.L., Packer, J.L., Arnold, W.A., McNeill, K., 2003b. Photochemical fate of pharmaceuticals in the environment: cimetidine and ranitidine. *Environ. Sci. Technol.* 37, 3342-3350. <https://doi.org/10.1021/es0340782>
- Le Page, G., Gunnarsson, L., Snape, J., Tyler, C.R., 2017. Integrating human and environmental health in antibiotic risk assessment: A critical analysis of protection goals, species sensitivity and antimicrobial resistance. *Environ. Int.* 109, 155–169. <https://doi.org/10.1016/j.envint.2017.09.013>
- Le Roux, D.M., Powers, L.C., Blough, N.V., 2021. Photoproduction rates of one-electron reductants by chromophoric dissolved organic matter via fluorescence spectroscopy: Comparison with superoxide and hydrogen peroxide rates. *Environ. Sci. Technol.* 55, 12095–12105. <https://doi.org/10.1021/acs.est.1c04043>
- Leavitt, P.R., Vinebrooke, R.D., Donald, D.B., Smol, J.P., Schindler, D.W., 1997. Past ultraviolet radiation environments in lakes derived from fossil pigments. *Nature* 388, 457–459. <https://doi.org/10.1038/41296>
- Levine, J.S., MacNichol, E.F. 1982. Color vision in fishes. *Scientific American*, 246, 140-149. <https://doi.org/10.1038/scientificamerican0282-140>
- Lewis, W.M., 1987. Tropical limnology. *Annu. Rev. Ecol. Syst.* 18, 159–184. <https://doi.org/10.1146/annurev.es.18.110187.001111>
- Li, C., Xie, H.B., Chen, J.W., Yang, X.H., Zhang, Y.F., Qiao, X.L., 2014. Predicting gaseous reaction rates of short chain chlorinated paraffins with ·OH: Overcoming the difficulty in experimental determination. *Environ. Sci. Technol.* 48, 13808-13816. <https://doi.org/10.1021/es504339r>

- Li, Y.J., Qiao, X.L., Zhang, Y.N., Zhou, C.Z., Xie, H.J., Chen, J.W., 2016. Effects of halide ions on photodegradation of sulfonamide antibiotics: formation of halogenated intermediates. *Water Research* 102, 405–412. <https://doi.org/10.1016/j.watres.2016.06.054>
- Lin, J., Chen, J.W., Wang, Y., Cai, X.Y., Wei, X.X., Qiao, X.L., 2008. More toxic and photoresistant products from photodegradation of Fenoxaprop-*p*-ethyl. *J. Agric. Food Chem.* 56, 8226–8230. <https://doi.org/10.1021/jf801341s>
- Liu, T., Yin, K., Liu, C., Luo, J., Crittenden, J., Zhang, W., Luo, S., He, Q., Deng, Y., Liu, H., Zhang, D., 2018. The role of reactive oxygen species and carbonate radical in oxcarbazepine degradation via UV, UV/H₂O₂. Kinetics, mechanisms and toxicity evaluation. *Water Res.* 147, 204–213. <https://doi.org/10.1016/j.watres.2018.10.007>
- Liu, Y., Mekic, M., Carena, L., Vione D., Gligorovski, S., Zhang, G., Jin, B., 2020. Tracking photodegradation products and bond-cleavage reaction pathways of triclosan using ultra-high resolution mass spectrometry and stable carbon isotope analysis. *Environ. Pollut.* 264, 114673. <https://doi.org/10.1016/j.envpol.2020.114673>
- Liu, Y.J., Yan, S.W., Lian, L.S., Ma, J.Z., Zhou, H.X., Song, W.H., 2019. Assessing the contribution of hydroxylation species in the photochemical transformation of primidone (pharmaceutical). *Sci. Total. Environ.* 696, 133826. <https://doi.org/10.1016/j.scitotenv.2019.133826>
- Lueder, U., Jørgensen, B.B., Kappler, A., Schmidt, C., 2020. Photochemistry of iron in aquatic environments. *Environ. Sci. Process. Impacts* 22, 12–24. <https://doi.org/10.1039/c9em00415g>
- Lund-Hansen, L.C., 2004. Diffuse attenuation coefficients $K_d(\text{PAR})$ at the estuarine North Sea-Baltic Sea transition: Time-series, partitioning, absorption, and scattering. *Estuar. Coast. Shelf Sci.* 61, 251–259. <https://doi.org/10.1016/j.ecss.2004.05.004>

- Luo, T.L., Chen, J.W., Li, X.H., Zhang, S.Y., Yao, H.Y., Peijnenburg, W.J.G.M., 2018. Effects of lomefloxacin on survival, growth and reproduction of *Daphnia magna* under simulated sunlight radiation. *Ecotoxicol. Environ. Saf.*, 166, 63–70. <https://doi.org/10.1016/j.ecoenv.2018.09.067>
- Luo, X., Wei, X.X., Chen, J.W., Xie, Q., Yang, X.H., Peijnenburg, W.J.G.M., 2019. Rate constants of hydroxyl radicals reaction with different dissociation species of fluoroquinolones and sulfonamides: Combined experimental and QSAR studies. *Water. Res.* 166, 115083. <https://doi.org/10.1016/j.watres.2019.115083>
- Luo, X., Yang, X.H., Qiao, X.L., Wang, Y., Chen, J.W., Wei, X.X., Peijnenburg, W.J.G.M., 2017. Development of a QSAR model for predicting aqueous reaction rate constants of organic chemicals with hydroxyl radicals. *Environ. Sci. Process. Impacts* 19, 350-356. <https://doi.org/10.1039/C6EM00707D>
- Ma, J.Z., Nie, J.X., Zhou, H.X., Wang, H., Lian, L.S., Yan, S.W., Song, W.H., 2020. Kinetic consideration of photochemical formation and decay of superoxide radical in dissolved organic matter solutions. *Environ. Sci. Technol.* 54, 3199–3208. <https://doi.org/10.1021/acs.est.9b06018>
- Mack, J., Bolton, J.R., 1999. Photochemistry of nitrite and nitrate in aqueous solution: A review. *J. Photochem. Photobiol. A Chem.* 128, 1–13. [https://doi.org/10.1016/S1010-6030\(99\)00155-0](https://doi.org/10.1016/S1010-6030(99)00155-0)
- Mackay, D., 2001. The fugacity approach, *Multimedia Environmental Models: The Fugacity Approach*, Second Edition. <https://doi.org/10.1201/9781420032543>
- Madikizela, L.M., Ncube, S., Chimuka, L., 2020. Analysis, occurrence and removal of pharmaceuticals in African water resources: A current status. *J. Environ. Manage.* 253, 109741. <https://doi.org/10.1016/j.jenvman.2019.109741>
- Margot, J., Kienle, C., Magnet, A., Weil, M., Rossi, L., de Alencastro, L.F., Abegglen, C., Thonney, D., Chèvre, N., Schärer, M., Barry, D.A., 2013. Treatment of micropollutants in municipal wastewater: Ozone or powdered activated carbon?

- Sci. Total Environ. 461–462, 480–498.
<https://doi.org/10.1016/j.scitotenv.2013.05.034>
- Markager, S., Vincent, W.F., 2000. Spectral light attenuation and the absorption of UV and blue light in natural waters. *Limnol. Oceanogr.* 45, 642–650.
<https://doi.org/10.4319/lo.2000.45.3.0642>
- Mathon, B., Coquery, M., Miège, C., Vandycke, A., Choubert, J.M., 2019. Influence of water depth and season on the photodegradation of micropollutants in a free-water surface constructed wetland receiving treated wastewater. *Chemosphere* 235, 260–270. <https://doi.org/10.1016/j.chemosphere.2019.06.140>
- Mazellier, P., Busset, C., Delmont, A., De Laat, J., 2007. A comparison of fenuron degradation by hydroxyl and carbonate radicals in aqueous solution. *Water Res.* 41, 4585–4594. <https://doi.org/10.1016/j.watres.2007.06.066>
- McKay, G., 2020. Emerging investigator series: Critical review of photophysical models for the optical and photochemical properties of dissolved organic matter. *Environ. Sci. Process. Impacts* 22, 1139–1165.
<https://doi.org/10.1039/d0em00056f>
- McKay, G., Couch, K.D., Mezyk, S.P., Rosario-Ortiz, F.L., 2016. Investigation of the coupled effects of molecular weight and charge-Transfer interactions on the optical and photochemical properties of dissolved organic matter. *Environ. Sci. Technol.* 50, 8093–8102. <https://doi.org/10.1021/acs.est.6b02109>
- McKay, G., Huang, W.X., Romera-Castillo, C., Crouch, J.E., Rosario-Ortiz, F.L., Jaffé, R., 2017. Predicting reactive intermediate quantum yields from dissolved organic matter photolysis using optical properties and antioxidant capacity. *Environ. Sci. Technol.* 51, 5404–5413. <https://doi.org/10.1021/acs.est.6b06372>
- McKay, G., Rosario-Ortiz, F.L., 2015. Temperature dependence of the photochemical formation of hydroxyl radical from dissolved organic matter. *Environ. Sci. Technol.* 49, 4147–4154. <https://doi.org/10.1021/acs.est.5b00102>

- McNeill, K., Canonica, S., 2016. Triplet state dissolved organic matter in aquatic photochemistry: Reaction mechanisms, substrate scope, and photophysical properties. *Environ. Sci. Process. Impacts* 18, 1381–1399. <https://doi.org/10.1039/c6em00408c>
- Michael, I., Achilleos, A., Lambropoulou, D., Torrens, V.O., Pérez, S., Petrović, M., Barceló, D., Fatta-Kassinos, D., 2014. Proposed transformation pathway and evolution profile of diclofenac and ibuprofen transformation products during (sono)photocatalysis. *Appl. Catal. B Environ.* 147, 1015–1027. <https://doi.org/10.1016/j.apcatb.2013.10.035>
- Minero, C., Chiron, S., Falletti, G., Maurino, V., Pelizzetti, E., Ajassa, R., Carlotti, M.E., Vione, D., 2007. Photochemical processes involving nitrite in surface water samples. *Aquat. Sci.* 69, 71–85. <https://doi.org/10.1007/s00027-007-0881-6>
- Moran, M.A., Sheldon, W.M., Zepp, R.G., 2000. Carbon loss and optical property changes during long-term photochemical and biological degradation of estuarine dissolved organic matter. *Limnol. Oceanogr.* 45, 1254–1264. <https://doi.org/10.4319/lo.2000.45.6.1254>
- Mossoba, M.M., Makino, K., Riesz, P., 1982. Photoionization of aromatic amino acids in aqueous solutions. A spin-trapping and electron spin resonance study. *J. Phys. Chem.* 86, 3478–3483. <https://doi.org/10.1021/j100214a041>
- Mulholland, P.J., Hill, W.R., 1997. Seasonal patterns in streamwater nutrient and dissolved organic carbon concentrations: Separating catchment flow path and in-stream effects. *Water Resour. Res.* 33, 1297–1306. <https://doi.org/10.1029/97WR00490>
- Musiolková, M., Huszár, P., Navrátil, M., Špunda, V., 2021. Impact of season, cloud cover, and air pollution on different spectral regions of ultraviolet and visible incident solar radiation at the surface. *Q. J. R. Meteorol. Soc.* 147, 2834–2849. <https://doi.org/10.1002/qj.4102>

- Myers, D.R., 2013. Introduction to solar radiation measurements, solar radiation. First edition. CRC Press. Boca Rato, USA. <https://doi.org/10.1201/b13898-2>
- Nebbioso, A., Piccolo, A., 2013. Molecular characterization of dissolved organic matter (DOM): A critical review. *Anal. Bioanal. Chem.* 405, 109–124. <https://doi.org/10.1007/s00216-012-6363-2>
- Niu, J.F., Huang, L.P., Chen, J.W., Yu, G., Schramm, K.-W., 2005. Quantitative structure–property relationships on photolysis of PCDD/Fs adsorbed to spruce (*Picea abies* (L.) Karst.) needle surfaces under sunlight irradiation. *Chemosphere* 58, 917–924. <https://doi.org/10.1016/j.chemosphere.2004.09.051>
- Niu, J.F., Zhang, L.L., Li, Y., Zhao, J.B., Lv, S.D., Xiao, K.Q., 2013. Effects of environmental factors on sulfamethoxazole photodegradation under simulated sunlight irradiation: Kinetics and mechanism. *J. Environ. Sci. (China)* 25, 1098–1106. [https://doi.org/10.1016/S1001-0742\(12\)60167-3](https://doi.org/10.1016/S1001-0742(12)60167-3)
- Niu, X.Z., Liu, C., Gutierrez, L., Croué, J.P., 2014. Photobleaching-induced changes in photosensitizing properties of dissolved organic matter. *Water Res.* 66, 140–148. <https://doi.org/10.1016/j.watres.2014.08.017>
- Ossola, R., Jönsson, O.M., Moor, K., McNeill, K., 2021. Singlet oxygen quantum yields in environmental waters. *Chem. Rev.* 121, 4100–4146. <https://doi.org/10.1021/acs.chemrev.0c00781>
- Ossola, R., Schmitt, M., Erickson, P.R., McNeill, K., 2019. Furan carboxamides as model compounds to study the competition between two modes of indirect photochemistry. *Environ. Sci. Technol.* 53, 9594–9603. <https://doi.org/10.1021/acs.est.9b02895>
- Page, S.E., Arnold, W.A., McNeill, K., 2011. Assessing the contribution of free hydroxyl radical in organic matter-sensitized photohydroxylation reactions. *Environ. Sci. Technol.* 45, 2818–2825. <https://doi.org/10.1021/es2000694>

- Parker, K.M., Mitch, W.A., 2016. Halogen radicals contribute to photooxidation in coastal and estuarine waters. *Proc. Natl. Acad. Sci. U. S. A.* 113, 5868–5873. <https://doi.org/10.1073/pnas.1602595113>
- Parker, K.M., Pignatello, J.J., Mitch, W.A., 2013. Influence of ionic strength on triplet-state natural organic matter loss by energy transfer and electron transfer pathways. *Environ. Sci. Technol.* 47, 10987-10944. <https://doi.org/10.1021/es401900j>
- Partanen, S.B., Apell, J.N., Lin, J., McNeill, K., 2021. Factors affecting the mixed-layer concentrations of singlet oxygen in sunlit lakes. *Environ. Sci. Process. Impacts* 23, 1130–1145. <https://doi.org/10.1039/d1em00062d>
- Patel, M., Kumar, R., Kishor, K., Mlsna, T., Pittman, C.U., Mohan, D., 2019. Pharmaceuticals of emerging concern in aquatic systems: Chemistry, occurrence, effects, and removal methods. *Chem. Rev.* 119, 3510–3673. <https://doi.org/10.1021/acs.chemrev.8b00299>
- Petasne, R.G., Zika, R.G., 1987. Fate of superoxide in coastal sea water. *Nature* 325, 516–518. <https://doi.org/10.1038/325516a0>
- Peterson, B.M., McNally, A.M., Cory, R.M., Thoemke, J.D., Cotner, J.B., McNeill, K., 2012. Spatial and temporal distribution of singlet oxygen in lake superior. *Environ. Sci. Technol.* 46, 7222–7229. <https://doi.org/10.1021/es301105e>
- Petrović, M., Škrbić, B., Živančev, J., Ferrando-Climent, L., Barcelo, D., 2014. Determination of 81 pharmaceutical drugs by high performance liquid chromatography coupled to mass spectrometry with hybrid triple quadrupole–linear ion trap in different types of water in Serbia. *Sci. Total. Environ.* 468–469, 415–428. <https://doi.org/10.1016/J.SCITOTENV.2013.08.079>
- Plumlee, M.H., McNeill, K., Reinhard, M., 2009. Indirect photolysis of perfluorochemicals: hydroxyl radical-initiated oxidation of N-ethyl perfluorooctane sulfonamido acetate (N-EtFOSAA) and other

- perfluoroalkanesulfonamides. *Environ. Sci. Technol.* 43, 3662-3668.
<https://doi.org/10.1021/es803411w>
- Rehrl, A.L., Golovko, O., Ahrens, L., Köhler, S., 2020. Spatial and seasonal trends of organic micropollutants in Sweden's most important drinking water reservoir. *Chemosphere* 249, 1–9. <https://doi.org/10.1016/j.chemosphere.2020.126168>
- Ribeiro, A.R., Nunes, O.C., Pereira, M.F.R., Silva, A.M.T., 2015. An overview on the advanced oxidation processes applied for the treatment of water pollutants defined in the recently launched Directive 2013/39/EU. *Environ. Int.* 75, 33–51.
<https://doi.org/10.1016/j.envint.2014.10.027>
- Robert, A.P., Alloy, M.M., Oris, J.T., 2017. Review of the photo-induced toxicity of environmental contaminants. *Comp. Biochem. Phys. C*, 191, 160-167.
<https://doi.org/10.1016/j.cbpc.2016.10.005>
- Roots, E.F., 1989. Climate change: High-latitude regions. *Clim. Change* 15, 223–253.
<https://doi.org/10.1007/BF00138853>
- Rose, K.C., Hamilton, D.P., Williamson, C.E., McBride, C.G., Fischer, J.M., Olson, M.H., Saros, J.E., Allan, M.G., Cabrol, N., 2014. Light attenuation characteristics of glacially-fed lakes. *J. Geophys. Res. Biogeosciences* 119, 1446–1457. <https://doi.org/10.1002/2014JG002674>
- Rossignol, S., Tinel, L., Bianco, A., Passananti, M., Brigante, M., Donaldson, D.J., George, C., 2016. Atmospheric photochemistry at a fatty acid-coated air-water interface. *Science* 353, 699–702. <https://doi.org/10.1126/science.aaf3617>
- Rozas, O., Baeza, C., Núñez, K., Rossner, A., Urrutia, R., Mansilla, H.D., 2017. Organic micropollutants (OMPs) oxidation by ozone: Effect of activated carbon on toxicity abatement. *Sci. Total Environ.* 590–591, 430–439.
<https://doi.org/10.1016/j.scitotenv.2016.12.120>
- Rubasinghe, G., Gurung, R., Rijal, H., Maldonado-Torres, S., Chan, A., Acharya, S., Rogelj, S., Piyasena, M., 2018. Abiotic degradation and environmental toxicity

- of ibuprofen: Roles of mineral particles and solar radiation. *Water Res.* 131, 22–32. <https://doi.org/10.1016/J.WATRES.2017.12.016>
- Sanganyado, E., Gwenzi, W., 2019. Antibiotic resistance in drinking water systems: Occurrence, removal, and human health risks. *Sci. Total. Environ.* 669, 785–797. <https://doi.org/10.1016/j.scitotenv.2019.03.162>
- Scheffer, M., 2004. *Ecology of shallow lakes*. Springer Dordrecht, Netherlands,. <https://doi.org/10.1007/978-1-4020-3154-0>
- Schmitt, M., Wack, K., Glaser, C., Wei, R., Zwiener, C., 2021. Separation of photochemical and non-photochemical diurnal in-stream attenuation of micropollutants. *Environ. Sci. Technol.* 55, 8908–8917. <https://doi.org/10.1021/acs.est.1c02116>
- Schwaiger, J., Ferling, H., Mallow, U., Wintermayr, H., Negele, R.D., 2004. Toxic effects of the non-steroidal anti-inflammatory drug diclofenac. Part I: Histopathological alterations and bioaccumulation in rainbow trout. *Aquat. Toxicol.* 68, 141–150. <https://doi.org/10.1016/j.aquatox.2004.03.014>
- Scully, N.M., McQueen, D.J., Lean, D.R.S., 1996. Hydrogen peroxide formation: The interaction of ultraviolet radiation and dissolved organic carbon in lake waters along a 43–75°N gradient. *Limnol. Oceanogr.* 41, 540–548. <https://doi.org/10.4319/lo.1996.41.3.0540>
- Shang, J., Chen, J., Shen, Z.Y., Xiao, X.Z, Yang, H.N., Wang, Y., Ruan, A.D., 2015. Photochemical degradation of PAHs in estuarine surface water: effects of DOM, salinity, and suspended particulate matter. *Environ. Sci. Pollut. Res.* 22, 12374–12383. <https://doi.org/10.1007/s11356-015-4543-2>
- Shao, J.P., Chen, J.W., Xie, Q., Wang, Y., Li, X.H., Hao, C., 2010. Electron-accepting potential of solvents determines photolysis rates of polycyclic aromatic hydrocarbons: Experimental and density functional theory study. *J. Hazard. Mater.* 179, 173–177. <https://doi.org/10.1016/j.jhazmat.2010.02.075>

- Sharpless, C.M., Blough, N.V., 2014. The importance of charge-transfer interactions in determining chromophoric dissolved organic matter (CDOM) optical and photochemical properties. *Environ. Sci. Process. Impacts* 16, 654–671. <https://doi.org/10.1039/c3em00573a>
- Shih, Y. Hsin, Wang, C.K., 2009. Photolytic degradation of polybromodiphenyl ethers under UV-lamp and solar irradiations. *J. Hazard. Mater.* 165, 34–38. <https://doi.org/10.1016/j.jhazmat.2008.09.103>
- Sommaruga, R., 2001. The role of solar UV radiation in the ecology of alpine lakes. *J. Photochem. Photobiol. B Biol.* 62, 35–42. [https://doi.org/10.1016/S1011-1344\(01\)00154-3](https://doi.org/10.1016/S1011-1344(01)00154-3)
- Soneye, O.O., Ayoola, M.A., Ajao, I.A., Jegede, O.O., 2019. Diurnal and seasonal variations of the incoming solar radiation flux at a tropical station, Ile-Ife, Nigeria. *Heliyon* 5, E01673. <https://doi.org/10.1016/j.heliyon.2019.e01673>
- Söregård, M., Campos-Pereira, H., Ullberg, M., Lai, F.Y., Golovko, O., Ahrens, L., 2019. Mass loads, source apportionment, and risk estimation of organic micropollutants from hospital and municipal wastewater in recipient catchments. *Chemosphere*. <https://doi.org/10.1016/j.chemosphere.2019.06.041>
- Southworth, B.A., Voelker, B.M., 2003. Hydroxyl radical production via the photo-fenton reaction in the presence of fulvic acid. *Environ. Sci. Technol.* 37, 1130–1136. <https://doi.org/10.1021/es020757l>
- Sulzberger, B., Canonica, S., Egli, T., Giger, W., Klausen, J., Von Gunten, U., 1997. Oxidative transformations of contaminants in natural and in technical systems. *Chimia*. 51, 900–907.
- Sun, L., Qian, J., Blough, N. V., Mopper, K., 2015. Insights into the photoproduction sites of hydroxyl radicals by dissolved organic matter in natural waters. *Environ. Sci. Technol. Lett.* 2, 352–356. <https://doi.org/10.1021/acs.estlett.5b00294>

- Sun, P., Pavlostathis, S.G., Huang, C.H., 2014. Photodegradation of veterinary ionophore antibiotics under UV and solar irradiation. *Environ. Sci. Technol.* 48, 13188–13196. <https://doi.org/10.1021/es5034525>
- Sun, Y.L., Xie, H.B., Zhou, C.Z, Wu, Y.D., Pu, M.J., Niu, J.F., 2020. The role of carbonate in sulfamethoxazole degradation by peroxymonosulfate without catalyst and the generation of carbonate radical. *J. Hazard. Mater.* 398, 122827. <https://doi.org/10.1016/j.jhazmat.2020.122827>
- Thirumavalavan, M., Hu, Y.L., Lee, J.F., 2012. Effects of humic acid and suspended solids on adsorption and photo-degradation of microcystin-LR onto samples from Taiwan reservoirs and rivers. *J. Hazard. Mater.* 217–218, 323–329. <https://doi.org/10.1016/j.jhazmat.2012.03.031>
- Valero, F.P.J., Minnis, P., Pope, S.K., Bucholtz, A., Bush, B.C., Doelling, D.R., Smith, W.L., Dong, X., 2000. Absorption of solar radiation by the atmosphere as determined using satellite, aircraft, and surface data during the Atmospheric Radiation Measurement Enhanced Shortwave Experiment (ARESE). *J. Geophys. Res. Atmos.* 105, 4743–4758. <https://doi.org/10.1029/1999JD901063>
- Vimalkumar, K., Arun, E., Krishna-Kumar, S., Poopal, R.K., Nikhil, N.P., Subramanian, A., Babu-Rajendran, R., 2018. Occurrence of triclocarban and benzotriazole ultraviolet stabilizers in water, sediment, and fish from Indian rivers. *Sci. Total Environ.* 625, 1351–1360. <https://doi.org/10.1016/J.SCITOTENV.2018.01.042>
- Vione, D., 2020. A critical view of the application of the APEX software (Aqueous Photochemistry of Environmentally-Occurring Xenobiotics) to predict photoreaction kinetics in surface freshwaters. *Molecules* 25. <https://doi.org/10.3390/molecules25010009>
- Vione, D. and Carena, L. 2020. The possible production of harmful intermediates is the “Dark Side” of the environmental photochemistry of contaminants

- (potentially adverse effects, and many knowledge gaps). *Environ. Sci. Technol.* 54, 5328-5330. <https://doi.org/10.1021/acs.est.0c01049>
- Vione, D., Encinas, A., Fabbri, D., Calza, P., 2018. A model assessment of the potential of river water to induce the photochemical attenuation of pharmaceuticals downstream of a wastewater treatment plant (Guadiana River, Badajoz, Spain). *Chemosphere* 198, 473–481. <https://doi.org/10.1016/j.chemosphere.2018.01.156>
- Vione, D., Minella, M., Maurino, V., Minero, C., 2014. Indirect photochemistry in sunlit surface waters: Photoinduced production of reactive transient species. *Chem. - A Eur. J.* 20, 10590–10606. <https://doi.org/10.1002/chem.201400413>
- Vione, D., Scozzaro, A., 2019. Photochemistry of surface fresh waters in the framework of climate change. *Environ. Sci. Technol.* 53, 7945–7963. <https://doi.org/10.1021/acs.est.9b00968>
- Wang, H., Zhou, H.X, Ma, J.Z, Nie, J.X, Yan, S.W, Song, W.H, 2020a. Triplet photochemistry of dissolved black carbon and its effects on the photochemical formation of reactive oxygen species. *Environ. Sci. Technol.* 54, 4903–4911. <https://doi.org/10.1021/acs.est.0c00061>
- Wang, J.Q, Chen, J.W, Qiao, X.L, Wang, Y., Cai, X.Y, Zhou, C.Z, Zhang, Y.L., Ding, G.H, 2018. DOM from mariculture ponds exhibits higher reactivity on photodegradation of sulfonamide antibiotics than from offshore seawaters. *Water Res.* 144, 365–372. <https://doi.org/10.1016/j.watres.2018.07.043>
- Wang, J.Q, Chen, J.W, Qiao, X.L, Zhang, Y.N, Uddin, M., Guo, Z.Y., 2019. Disparate effects of DOM extracted from coastal seawaters and freshwaters on photodegradation of 2,4-Dihydroxybenzophenone. *Water Res.* 151, 280–287. <https://doi.org/10.1016/j.watres.2018.12.045>
- Wang, J.Q, Wang, K., Guo, Y.C, Niu, J.F, 2020b. Photochemical degradation of neбиволol in different natural organic matter solutions under simulated sunlight

- irradiation: Kinetics, mechanism and degradation pathway. *Water Res.* 173, 115524. <https://doi.org/10.1016/j.watres.2020.115524>
- Wang, J.Q., Wang, K., Guo, Y.C., Ye, Z.M., Guo, Z.Y., Lei, Y., Yang, X., Zhang, L.L., Niu, J.F., 2021a. Dichlorine radicals (Cl_2^{\bullet}) promote the photodegradation of propranolol in estuarine and coastal waters. *J. Hazard. Mater.* 414, 125536. <https://doi.org/10.1016/j.jhazmat.2021.125536>
- Wang, Y., Chen, J.W., Ge, L.K., Wang, D.G., Cai, X.Y., Huang, L.P., Hao, C. 2009. Experimental and theoretical studies on the photoinduced acute toxicity of a series of anthraquinone derivatives towards the water flea (*Daphnia magna*). *Dyes Pigments*, 83, 276–280. <https://doi.org/10.1016/j.dyepig.2009.04.015>
- Wang, Y.F., Roddick, F.A., Fan, L.H., 2017. Direct and indirect photolysis of seven micropollutants in secondary effluent from a wastewater lagoon. *Chemosphere* 185, 297–308. <https://doi.org/10.1016/j.chemosphere.2017.06.122>
- Wang, Y.K., Ma, X.Y., Zhang, S., Tang, L., Zhang, H., Wang, X.C., 2021b. Sunlight-induced changes in naturally stored reclaimed water: Dissolved organic matter, micropollutant, and ecotoxicity. *Sci. Total Environ.* 753, 141768. <https://doi.org/10.1016/j.scitotenv.2020.141768>
- Wardman, P., 1989. Reduction potentials of one electron couples involving free radicals in aqueous solution. *J. Phys. Chem. Ref. Data* 18, 1637–1755. <https://doi.org/10.1063/1.555843>
- Wayne, R.P., 2005. Basic concepts of photochemical transformations. *Environ. Photochem. Part II*, 1–47. <https://doi.org/10.1007/b138178>
- Wei, X.X., Chen, J.W., Xie, Q., Zhang, S.Y., Ge, L.K., Qiao, X.L., 2013. Distinct photolytic mechanisms and products for different dissociation species of ciprofloxacin. *Environ. Sci. Technol.* 47, 4284–4290. <https://doi.org/10.1021/es400425b>
- Wei, X.X., Chen, J.W., Xie, Q., Zhang, S.Y., Li, Y.J., Zhang, Y.F., Xie, H.B., 2015. Photochemical behavior of antibiotics impacted by complexation effects of

- concomitant metals: a case for ciprofloxacin and Cu(II). 17, 1220-1227.
<https://doi.org/10.1039/C5EM00204D>
- Wenk, J., Canonica, S., 2012. Phenolic antioxidants inhibit the triplet-induced transformation of anilines and sulfonamide antibiotics in aqueous solution. *Environ. Sci. Technol.* 46, 5455–5462. <https://doi.org/10.1021/es300485u>
- Werner, J.J., Arnold, W.A., McNeill, K., 2006. Water hardness as a photochemical parameter: tetracycline photolysis as a function of calcium concentration, magnesium concentration, and pH. *Environ. Sci. Technol.* 40, 7236–7241.
<https://doi.org/10.1021/es060337m>
- Weyhenmeyer, G.A., Müller, R.A., Norman, M., Tranvik, L.J., 2016. Sensitivity of freshwaters to browning in response to future climate change. *Clim. Change* 134, 225–239. <https://doi.org/10.1007/s10584-015-1514-z>
- White, E.M., Vaughan, P.P., Zepp, R.G., 2003. Role of the photo-Fenton reaction in the production of hydroxyl radicals and photobleaching of colored dissolved organic matter in a coastal river of the southeastern United States. *Aquat. Sci.* 65, 402-414. <https://doi.org/10.1007/s00027-003-0675-4>
- Wu, B.B., Liu, T., Wang, Y.L., Zhao, G.Q., Chen, B.L., Chu, C.H., 2021. High sample throughput LED reactor for facile characterization of the quantum yield spectrum of photochemically produced reactive intermediates. *Environ. Sci. Technol.* 55, 16204-16214. <https://doi.org/10.1021/acs.est.1c04608>
- Wu, C., Linden, K.G., 2010. Phototransformation of selected organophosphorus pesticides: Roles of hydroxyl and carbonate radicals. *Water Res.* 44, 3585–3594.
<https://doi.org/10.1016/j.watres.2010.04.011>
- Wu, Y., Bu, L., Duan, X., Zhu, S., Kong, M., Zhu, N., Zhou, S., 2020. Mini review on the roles of nitrate/nitrite in advanced oxidation processes: Radicals transformation and products formation. *J. Clean. Prod.* 273, 123065.
<https://doi.org/10.1016/j.jclepro.2020.123065>

- Xu, T., Chen, J.W., Chen, X., Xie, H.J., Wang, Z.Y., Xia, D.M., Tang, W.H., Xie, H.B., 2021. Prediction models on pKa and base-catalyzed hydrolysis kinetics of parabens: Experimental and quantum chemical studies. *Environ. Sci. Technol.* 55, 6022–6031. <https://doi.org/10.1021/acs.est.0c06891>
- Xu, T., Chen, J.W., Wang, Z.Y., Tang, W.H., Xia, D.M., Fu, Z.Q., Xie, H.J., 2019. Development of prediction models on base-catalyzed hydrolysis kinetics of phthalate esters with density functional theory calculation. *Environ. Sci. Technol.* 53, 5828–5837. <https://doi.org/10.1021/acs.est.9b00574>
- Yan, S.W., Liu, Y.J., Lian, L.S., Li, R., Ma, J.Z., Zhou, H.X., Song, W.H., 2019. Photochemical formation of carbonate radical and its reaction with dissolved organic matters. *Water Res.* 161, 288–296. <https://doi.org/10.1016/j.watres.2019.06.002>
- Yan, S.W., Song, W.H., 2014. Photo-transformation of pharmaceutically active compounds in the aqueous environment: A review. *Environ. Sci. Process. Impacts* 16, 697–720. <https://doi.org/10.1039/c3em00502j>
- Yan, S.W., Sun, J.Q., Sha, H.T., Li, Q., Nie, J.X., Zou, J.M., Chu, C.H., Song, W.H., 2021. Microheterogeneous distribution of hydroxyl radicals in illuminated dissolved organic matter solutions. *Environ. Sci. Technol.* 55, 10524–10533. <https://doi.org/10.1021/acs.est.1c03346>
- Yang, C., Song, G., Lim, W., 2020. A review of the toxicity in fish exposed to antibiotics. *Comp. Biochem. Physiol. C*, 237, 108840. <https://doi.org/10.1016/j.cbpc.2020.108840>
- Yang, P., Chen, J.W., Chen, S., Yuan, X., Schramm, K.-W., Kettrup, A., 2003. QSPR models for physicochemical properties of polychlorinated diphenyl ethers. *Sci. Total. Environ.* 305, 65-76. [https://doi.org/10.1016/S0048-9697\(02\)00467-9](https://doi.org/10.1016/S0048-9697(02)00467-9)
- Yang, Y., Pignatello, J.J., 2017. Participation of the halogens in photochemical reactions in natural and treated waters. *Molecules* 22. <https://doi.org/10.3390/molecules22101684>

- Yao, H., Sun, P.Z., Minakata, D., Crittenden, J.C., Huang, C.H., 2013. Kinetics and modeling of degradation of ionophore antibiotics by UV and UV/H₂O₂. *Environ. Sci. Technol.* 47, 4581–4589. <https://doi.org/10.1021/es3052685>
- Zareitalabad, P., Siemens, J., Hamer, M., Amelung, W., 2013. Perfluorooctanoic acid (PFOA) and perfluorooctanesulfonic acid (PFOS) in surface waters, sediments, soils and wastewater - A review on concentrations and distribution coefficients. *Chemosphere* 91, 725–732. <https://doi.org/10.1016/j.chemosphere.2013.02.02>
- Zepp, R.G., Braun, A.M., Holane, J., Leenheer, J.A., 1987. Photoproduction of hydrated electrons from natural organic solutes in aquatic environments. *Environ. Sci. Technol.* 21, 485–490. <https://doi.org/10.1021/es00159a010>
- Zepp, R.G., Cline, D.M., 1977. Rates of direct photolysis in aquatic environment. *Environ. Sci. Technol.* 11, 359–366. <https://doi.org/10.1021/es60127a013>
- Zepp, R.G., Faust, B.C., Holgne, J., 1992. Hydroxyl radical formation in aqueous reactions (pH 3-8) of iron(II) with hydrogen peroxide: the photo-Fenton reaction. *Environ. Sci. Technol.* 26, 313-319. <https://doi.org/10.1021/es00026a011>
- Zepp, R.G., Schlotzhauer, P.F., Sink, R.M., 1985. Photosensitized transformations involving electronic energy transfer in natural waters: Role of humic substances. *Environ. Sci. Technol.* 19, 74–81. <https://doi.org/10.1021/es00131a008>
- Zepp, R.G., Wolfe, N.L., Baughman, G.L., Hollis, R.C., 1977. Singlet oxygen in natural waters. *Nature* 267, 421–423. <https://doi.org/10.1038/267421a0>
- Zhang, K., Parker, K.M., 2018. Halogen radical oxidants in natural and engineered aquatic systems. *Environ. Sci. Technol.* 52, 9579–9594. <https://doi.org/10.1021/acs.est.8b02219>
- Zhang, Y., Del Vecchio, R., Blough, N. V., 2012. Investigating the mechanism of hydrogen peroxide photoproduction by humic substances. *Environ. Sci. Technol.* 46, 11836–11843. <https://doi.org/10.1021/es3029582>
- Zhang, Y.L., Green, N.W., Perdue, E.M., 2013. Acid-base properties of dissolved organic matter from pristine and oil-impacted marshes of Barataria Bay,

- Louisiana. Mar. Chem. 155, 42–49.
<https://doi.org/10.1016/j.marchem.2013.05.010>
- Zhang, Y.N., Wang, J.Q., Chen, J.W., Zhou, C.Z., Xie, Q., 2018a. Phototransformation of 2,3-Dibromopropyl-2,4,6-tribromophenyl ether (DPTE) in natural waters: Important roles of dissolved organic matter and chloride ion. Environ. Sci. Technol. 52, 10490–10499.
<https://doi.org/10.1021/acs.est.8b03258>
- Zhang, Y.N., Zhou, Y.J., Qu, J., Chen, J.W., Zhao, J.C., Lu, Y., Li, C., Xie, Q., Peijnenburg, W.J.G.M., 2018b. Unveiling the important roles of coexisting contaminants on photochemical transformations of pharmaceuticals: Fibrate drugs as a case study. J. Hazard. Mater. 358, 216–221.
<https://doi.org/10.1016/j.jhazmat.2018.06.068>
- Zhang, Y.Q., Zhang, J.F., Xiao, Y.J., Chang, V.W.C., Lim, T.T., 2017. Direct and indirect photodegradation pathways of cytostatic drugs under UV germicidal irradiation: Process kinetics and influences of water matrix species and oxidant dosing. J. Hazard. Mater. 324, 481–488.
<https://doi.org/10.1016/j.jhazmat.2016.11.016>
- Zhou, C.Z., Chen, J.W., Xie, H.J., Zhang, Y.N., Li, Y.J., Wang, Y., Xie, Q., Zhang, S.Y., 2018. Modeling photodegradation kinetics of organic micropollutants in water bodies: A case of the Yellow River estuary. J. Hazard. Mater. 349, 60–67.
<https://doi.org/10.1016/j.jhazmat.2018.01.051>
- Zhou, H.X., Yan, S.W., Lian, L.S., Song, W.H., 2019a. Triplet-State Photochemistry of Dissolved Organic Matter: Triplet-State Energy Distribution and Surface Electric Charge Conditions. Environ. Sci. Technol. 53, 2482–2490.
<https://doi.org/10.1021/acs.est.8b06574>
- Zhou, Y.J., Chen, C.Y., Guo, K.H., Wu, Z.H., Wang, L.P., Hua, Z.C., Fang, J.Y., 2020. Kinetics and pathways of the degradation of PPCPs by carbonate radicals in

advanced oxidation processes. *Water Res.* 185, 116231.

<https://doi.org/10.1016/j.watres.2020.116231>

Zhou, Y.J., Zhao, J.C., Zhang, Y.N., Qu, J., Li, C., Qin, W.C., Zhao, Y.H., Chen, J.W., Peijnenburg, W.J.G.M., 2019b. Trace amounts of fenofibrate acid sensitize the photodegradation of bezafibrate in effluents: Mechanisms, degradation pathways, and toxicity evaluation. *Chemosphere* 235, 900–907.

<https://doi.org/10.1016/J.CHEMOSPHERE.2019.07.008>

Zhu, Y., Kieber, D.J., 2018. Wavelength- and temperature-dependent apparent quantum yields for photochemical production of carbonyl compounds in the North Pacific Ocean. *Environ. Sci. Technol.* 52, 1929–1939.

<https://doi.org/10.1021/acs.est.7b05462>




Figure 1. Comparison and cumulative probability distributions of photodegradation and biodegradation (in fresh and sea waters) half-lives of commonly detected PPCPs, adapted from Baena-Nogueras et al. (2017)




Figure 2. Overview of photodegradation in aquatic environment.




Figure 3. Production of photochemically produced reactive intermediates in DOM solution. (k_d^T is relaxation rate constant of ${}^3\text{DOM}^*$ by O_2 -independent way; f_Δ is the proportion of ${}^3\text{DOM}^*$ reacting with O_2 that produces ${}^1\text{O}_2$; k_d^Δ is relaxation rate constant from ${}^1\text{O}_2$ to O_2) (Sharpless and Blough, 2014; McNeill and Canonica, 2016; Vione et al., 2014).




Figure 4. Variation of annual irradiance with latitude. [Data was collected from Lewis (1987)].




Figure 5. A) Diurnal and seasonal variation of the incoming solar radiation flux (SWdn) at Ile-Ife, Nigeria for the wet (March–October) and dry (November–February) seasons, during period of January 2016 to December 2017 (Soneye et al., 2019). B) Diurnal solar radiation variation (referred to a cloudless day October 22, 2015, in Dalian, 38.9° N latitude, range 280 nm – 500 nm) (Zhou et al., 2018).